



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Somato-dendritic vasopressin and oxytocin secretion in endocrine and autonomic regulation

Citation for published version:

Brown, CH, Ludwig, M, Tasker, JG & Stern, JE 2020, 'Somato-dendritic vasopressin and oxytocin secretion in endocrine and autonomic regulation', *Journal of Endocrinology*. <https://doi.org/oi.org/10.1111/jne.12856>

Digital Object Identifier (DOI):

[oi.org/10.1111/jne.12856](https://doi.org/10.1111/jne.12856)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Endocrinology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Journal of Neuroendocrinology**Somato-dendritic vasopressin and oxytocin secretion in endocrine and autonomic regulation**

Journal:	<i>Journal of Neuroendocrinology</i>
Manuscript ID	JNE-20-0008-RA.R1
Manuscript Type:	Review Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Brown, Colin; University of Otago, Department of Physiology Ludwig, Mike; University of Edinburgh, Centre for Discovery Brain Sciences Tasker, Jeffrey G.; Tulane University, Cell and Molecular Biology Stern, Javier; Georgia State University, Neuroscience
Keywords:	Somato-dendritic secretion, Paraventricular nucleus, Supraoptic nucleus, Oxytocin, Vasopressin

SCHOLARONE™
Manuscripts

Somato-dendritic vasopressin and oxytocin secretion in endocrine and autonomic regulation

Colin H. Brown ¹, Mike Ludwig ^{2,3}, Jeffrey G. Tasker ⁴ and Javier E. Stern ⁵

*¹ Brain Health Research Centre, Centre for Neuroendocrinology and Department of Physiology,
University of Otago, Dunedin, New Zealand*

²Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh, UK

³ Centre for Neuroendocrinology, Department of Immunology, University of Pretoria, Pretoria, South Africa

⁴ *Brain Institute and Department of Cell and Molecular Biology, Tulane University, New Orleans, LA, USA*

⁵Neuroscience Institute, Georgia State University, Atlanta, GA, USA

Running title: Somato-dendritic VP and OT secretion

Key words: Somato-dendritic secretion, paraventricular nucleus, supraoptic nucleus, oxytocin, vasopressin

Corresponding author: Colin H. Brown, Department of Physiology, University of Otago, Dunedin

9054, New Zealand. Tel.: +64-3-479-7354, Fax.: +64-3-479-7323, E-mail: colin.brown@otago.ac.nz

Abstract

Somato-dendritic secretion was first demonstrated over 30 years ago. However, while its existence has become widely accepted, the function of somato-dendritic secretion is still not completely understood. Hypothalamic magnocellular neurosecretory cells (MNCs) were among the first neuronal phenotypes in which somato-dendritic secretion was demonstrated and are among the neurones for which the functions of somato-dendritic secretion are best characterised. These neurones secrete the neuropeptides, vasopressin and oxytocin, in an orthograde manner from their axons in the posterior pituitary gland into the blood circulation to regulate body fluid balance and reproductive physiology. Retrograde somato-dendritic secretion of vasopressin and oxytocin modulate the activity of the neurones from which they are secreted, as well as the activity of neighbouring populations of neurones, to provide intra- and inter-population signals that coordinate the endocrine and autonomic responses for control of peripheral physiology. Somato-dendritic vasopressin and oxytocin have also been proposed to act as hormone-like signals in the brain. There is some evidence that somato-dendritic secretion from MNCs modulates the activity of neurones beyond their local environment where there are no vasopressin- or oxytocin-containing axons but, to date, there is no conclusive evidence for, or against, hormone-like signalling throughout the brain, although it is difficult to imagine that the levels of vasopressin found throughout the brain could be underpinned by release from relatively sparse axon terminal fields; the generation of data to resolve this issue remains a priority for the field.

Information transfer in the central nervous system

The classical understanding of communication in the nervous system is of synaptic transmission in a unidirectional manner within networks from presynaptic neurones to postsynaptic neurones. However, it has become clear that information transfer in the central nervous system is more complex than simple point-to-point, unidirectional transmission between neurones at synapses. Among the additional mechanisms that contribute to information transfer in the nervous system is somato-

dendritic secretion. Unlike classical synaptic transmission by neurotransmitters such as glutamate and GABA, which signals between pre- and postsynaptic neurones with spatial precision and high temporal resolution, somato-dendritic secretion causes longer-term changes than synaptic transmission that alters the overall excitability of neurones by modulating the strength of synaptic inputs and/or by modulating the baseline membrane potential. These effects can be autocrine or paracrine, on the neurone from which somato-dendritic secretion occurs or on nearby neurones, and might spread over relatively long distances to modulate the activity of neurones in brain areas distant from the site of secretion.

Somato-dendritic secretion occurs in many types of neurone and can involve many types of transmitter molecule (1). Magnocellular neurosecretory cells (MNCs) of the hypothalamic supraoptic nucleus (SON) and paraventricular nucleus (PVN) are among those for which the mechanisms and consequences of somato-dendritic secretion are best characterised. This review focusses on studies from the authors' laboratories, some of which were presented at the 22nd International Symposium on Regulatory Peptides, which have contributed to our understanding of how somato-dendritic secretion from MNCs contributes to endocrine and autonomic regulation of peripheral physiology in health and disease.

The magnocellular neurosecretory system

The magnocellular neurosecretory system comprises MNCs that predominantly secrete either vasopressin (the antidiuretic hormone) or oxytocin into the general circulation from the posterior pituitary gland (neurohypophysis). The principal function of vasopressin is to maintain body fluid balance and blood pressure by activation of renal V_2 -receptors to increase water reabsorption from the urine and, when blood pressure/volume is decreased, by activation of vascular V_{1a} -receptors ($V_{1a}Rs$) to cause vasoconstriction (2). The best-characterised physiological functions of oxytocin are to trigger uterine contractions during birth and milk ejection during lactation (2). However, oxytocin also contributes to body fluid balance by promoting natriuresis in the kidney (3) and by stimulating atrial natriuretic peptide secretion (4).

1
2
3 73 The human hypothalamus contains over 100,000 MNCs (5), with ~10,000 in the rat, that are
4
5 74 principally located in the SON and PVN as well as in several accessory nuclei (6). MNCs each project
6
7 75 a single axon to the posterior pituitary gland (Figure 1) and each axon branches extensively to form
8
9 76 several thousand neurosecretory axon swellings and terminals (7) that are each tightly packed with
10
11 77 dense core vesicles containing ~85,000 molecules of vasopressin or oxytocin in rats (8). Hormone
12
13 78 secretion is triggered by action potential invasion of the neurosecretory swellings and terminals. It has
14
15 79 been estimated that each MNC contains about 10 million dense core vesicles and secretes between
16
17 80 100 and 10,000 dense core vesicles from the posterior pituitary gland every minute to maintain basal
18
19 81 hormone concentrations in the circulation (9). Hence, the sustained output of the hormones, and the
20
21 82 consequent regulation of peripheral physiology, depends on the average action potential discharge
22
23 83 from the population (2, 10).
24
25
26
27 84 MNCs also synthesise lesser amounts of other neurotransmitters and neuromodulators that can be
28
29 85 contained in the same dense core vesicles as vasopressin or oxytocin (11), as well as glutamate-
30
31 86 containing microvesicles (12). To date, the only evidence of effects of these other neurotransmitters
32
33 87 and neuromodulators on peripheral physiology is for secretin, which increases renal antidiuresis (13).
34
35 88 Rather, their principal function appears to be modulation of hormone secretion at the level of the
36
37 89 posterior pituitary gland, which is comprehensively reviewed elsewhere (14), and at the level of the
38
39 90 somata and dendrites, as we describe here.
40
41
42
43 91 Some MNCs project axon collaterals to other brain areas. Originally, these were thought to remain
44
45 92 proximal to the SON (15) and PVN (16), projecting to local interneurons as part of a proposed local
46
47 93 feedback loop. More recently, it was shown that some MNC axon collaterals project more broadly
48
49 94 throughout the brain, with oxytocin MNCs projecting to the medial amygdala (MeA), central
50
51 95 amygdala (CeA), nucleus accumbens (17) and the lateral septum (18), and vasopressin MNCs to the
52
53 96 medial and lateral preoptic area, suprachiasmatic nucleus, lateral habenula, CeA, MeA (19, 20), locus
54
55 97 coeruleus (21) and arcuate nucleus (ARC) (22). These axon collaterals have been implicated in the
56
57 98 modulation of different behaviours, but it remains to be established how secretion from axon
58
59
60

collaterals to modulate behaviour relates to secretion from the posterior pituitary gland to modulate peripheral physiology.

MNCs possess 1 – 3 thick, varicose, aspiny dendrites of a few hundred micrometres in length. MNCs of the SON extend their dendrites to the ventral surface of the nucleus, where the dendrites bundle together within the ventral glial lamina (a layer of astrocytes on the ventral surface of the brain within the SON) (23) and MNCs of the PVN extend their dendrites towards the subependymal region of the third ventricle (24). In addition to being the site of afferent synaptic input, MNC dendrites are active players in shaping MNC activity through exocytosis of vasopressin and oxytocin (as well as other neurotransmitters/neuromodulators) into the extracellular space of the SON and PVN.

Somato-dendritic secretion from magnocellular neurosecretory cells

The somata and dendrites of MNCs are tightly packed with dense core vesicles containing either vasopressin or oxytocin (Figure 2), which undergo exocytosis to secrete their major neuropeptides (25, 26) along with lesser amounts of other co-packaged neurotransmitters and neuromodulators (11). Tannic acid capture of somato-dendritic secretion reveals that the entire vesicle content is released from MNCs (25).

Unlike synaptic transmission by classical neurotransmitters, dense core vesicle exocytosis from the somata and dendrites of MNCs requires a sustained increase in intracellular calcium (27, 28,) and calcium buffering limits increases in cytoplasmic calcium to restrain the activation of somato-dendritic secretion from MNCs (29). MNCs express different arrays of voltage-gated calcium channels in their somata and axon terminals (30). Relative to other voltage-gated calcium channels, N-type calcium channels ($Ca_{v2.2}$) carry a comparatively small current in MNC somata, but nevertheless contribute most significantly to somato-dendritic oxytocin secretion (31). While the primary trigger for somato-dendritic secretion is the influx of extracellular calcium, intracellular calcium release also contributes to somato-dendritic secretion from MNCs (27, 28).

1
2
3 123 Action potential invasion triggers exocytosis from axon terminals and MNC dendrites and appears to
4
5 124 support depolarisation-induced calcium spikes (32). Capacitance measurements from isolated MNCs
6
7 125 suggest that single action potentials trigger somato-dendritic secretion (33). However, functional
8
9 126 studies suggest that sustained intracellular calcium release is required to trigger somato-dendritic
10
11 127 secretion (27, 28). Furthermore, if every action potential fired by each MNC triggered somato-
12
13 128 dendritic secretion of a single dense core vesicle, the brain would be awash with vasopressin and
14
15 129 oxytocin (34). Exocytosis of ~6,000 dense core vesicles per second has been calculated to be
16
17 130 sufficient to maintain the concentrations of vasopressin and oxytocin measured in the rat
18
19 131 hypothalamus (34). There are ~10,000 MNCs in the rat hypothalamus (6). While ~25% of MNCs are
20
21 132 silent under basal conditions, active MNCs display a mean firing rate of ~5 Hz under basal conditions
22
23 133 (35). Hence, up to 37,500 action potentials are fired by MNCs every second, which is almost 10-fold
24
25 134 more than the number of dense core vesicles secreted. Hence, it is likely that trains of action potentials
26
27 135 that cause a more sustained depolarisation and calcium influx are required to trigger somato-dendritic
28
29 136 secretion from MNCs. Indeed, under basal conditions, some stimuli reduce the oxytocin MNC action
30
31 137 potential firing rate but increase somato-dendritic oxytocin secretion (36, 37), and it was shown
32
33 138 recently that action potential firing alone at physiological firing rates is insufficient to trigger
34
35 139 measurable somato-dendritic secretion from individual vasopressin MNCs (38).

36
37
38
39
40 140 In addition to permeation through voltage-gated calcium channels, calcium influx also occurs through
41
42 141 N-methyl-D-aspartate (NMDA) receptors (NMDARs), and synaptic NMDA receptors would be
43
44 142 expected to further increase cytoplasmic calcium concentrations during action potential firing.
45
46 143 Furthermore, MNCs express extrasynaptic NMDARs (39-41). While these extrasynaptic NMDARs
47
48 144 are activated by basal glutamate levels *in vitro* (39), they are not activated under basal conditions *in*
49
50 145 *vivo*, but are activated under stimulated conditions (35) and trigger somato-dendritic peptide secretion
51
52 146 (38).

53
54
55 147 In addition to triggering somato-dendritic secretion, increased cytoplasmic calcium also promotes
56
57 148 movement of dense core vesicles from the reserve pool toward the cell surface (42), where they are
58
59
60

ready for secretion in response to subsequent stimuli that raise cytoplasmic calcium. In parallel, increased intracellular calcium also promotes recruitment of N-type calcium channels (31) to make the system more sensitive to subsequent cytoplasmic calcium increases. Hence, this 'priming' increases somato-dendritic secretion triggered by subsequent signals that increase cytoplasmic calcium.

Action potential-mediated depolarisation is not the only trigger for somato-dendritic secretion from MNCs. Vasopressin and oxytocin MNCs express their respective receptors (43, 44) and activation of these receptors increases cytoplasmic calcium concentrations (45) to trigger somato-dendritic secretion (27, 28). While vasopressin and oxytocin trigger somato-dendritic secretion from vasopressin and oxytocin MNCs without a prior stimulus to prime the system, once the MNCs are primed, the peptides can trigger a much greater somato-dendritic secretion (27, 28).

There is an elaborate network of actin and microtubules in MNC somata and dendrites (46, 47). Cortical F-actin regulates somato-dendritic exocytosis; F-actin polymerization inhibits and F-actin depolymerisation increases somato-dendritic secretion from MNCs (48). Presumably, F-actin depolymerisation increases access to the plasma membrane and this process might account for the requirement for a sustained increase in intracellular calcium to trigger somato-dendritic secretion, since calcium causes F-actin depolymerisation. Unlike synaptic transmission, there does not appear to be any specific structure on the soma or dendrites that is specialised for somato-dendritic secretion (49), although it remains to be determined whether there are regions of the cortical F-actin network that are more readily depolymerised to allow dense core vesicles preferential access to the plasma membrane at specific sites for secretion.

Exocytosis can occur once the dense core vesicles reach the plasma membrane, which requires exocytotic machinery. The involvement of the soluble N-ethylmaleimide-sensitive factor attachment receptor (SNARE) complex in exocytosis from axon terminals is well established (50). While less well characterised, it appears that somato-dendritic secretion is also mediated by SNARE proteins. MNCs express SNARE proteins (51), however, while vesicle-associated membrane protein-2

(VAMP-2) and synaptosomal-associated protein 25 (SNAP-25) are both expressed in the axon terminals (52, 53), they are not expressed in the somata or dendrites of MNCs (54). Hence, the suite of SNARE complex proteins for somato-dendritic exocytosis likely differs from that for axon terminal secretion from MNCs.

179 **Stimulation of somato-dendritic secretion by neurotransmitters, peptides and hormones**

180 Noradrenergic afferents from the ventrolateral medulla (VLM) A1 cell group and the nucleus of the
181 tractus solitarius (NTS) A2 cell group make prominent projections to MNCs (2). NTS noradrenergic
182 afferents are activated during birth and lactation (55) as part of the Ferguson reflex (56), and
183 noradrenaline facilitates somato-dendritic oxytocin secretion in late pregnancy and lactation (57-59).
184 Oxytocin also increases noradrenaline secretion within the SON (60), which presumably establishes a
185 local positive feedback loop that reinforces oxytocin MNC excitation and promotes oxytocin secretion
186 into the circulation to trigger uterine contractions during birth and milk ejection during lactation
187 (Figure 3).

188 Proopiomelanocortin (POMC) afferents from the ARC project to the SON and PVN, particularly to
189 regions of the SON and PVN that are enriched in oxytocin MNCs (61). POMC neurones secrete α -
190 melanocyte stimulating hormone (α -MSH), which acts on melanocortin-4 (MC4-R) receptors in the
191 SON and PVN (62). While α -MSH inhibits oxytocin secretion into the circulation, it increases
192 somato-dendritic oxytocin secretion (37, 63). MC4-R activation increases intracellular calcium in
193 oxytocin MNCs to trigger somato-dendritic oxytocin secretion as well as endocannabinoid secretion,
194 which inhibits the activity of the MNCs to reduce axonal oxytocin secretion into the blood (37, 63).
195 Remarkably, α -MSH inhibition of oxytocin MNC activity is lost in mid-pregnancy (62), but it has yet
196 to be determined whether this represents a switch from pre-pregnancy inhibition to stimulation during
197 lactation, as is seen for prolactin effects on oxytocin MNCs (64). Furthermore, it is not known
198 whether α -MSH effects on somato-dendritic oxytocin secretion change in pregnancy and lactation.

199 In addition to neurotransmitters, other hormones can also trigger somato-dendritic secretion from
200 MNCs. The orexigenic hormone, ghrelin is synthesised by oxyntic cells in the gastric mucosa, but not
201 in the brain (65). Central ghrelin administration increases vasopressin secretion into the circulation via
202 activation of neuropeptide Y neurones (66). In addition, ghrelin stimulates somato-dendritic
203 vasopressin secretion, which increases adenosine triphosphate (ATP) release from astrocytes to
204 increase presynaptic GABA release onto the vasopressin MNCs (67) (Figure 4).

205 Autocrine/paracrine modulation of vasopressin magnocellular neurosecretory cell activity

206 The effects of vasopressin, oxytocin and other neurotransmitters and neuromodulators secreted from
207 the somata and dendrites of MNCs can be broadly categorised as autocrine, regulating the activity of
208 the MNC from which secretion occurs, and paracrine, regulating the activity of neighbouring neurons,
209 including other neuronal populations.

210 $V_{1a}R$ and V_{1b} receptors ($V_{1b}R$) are expressed in the membranes of vasopressin-containing dense core
211 vesicles (68) and are presumably inserted into the plasma membrane during somato-dendritic
212 vasopressin secretion. Hence, vasopressin receptors newly trafficked to the plasma membrane will be
213 exposed to high concentrations of vasopressin to underpin activity-dependent autocrine feedback
214 regulation of vasopressin MNC activity.

215 Vasopressin MNCs express a range of activity patterns under basal conditions; some are silent
216 throughout recordings, some display irregular activity, some are continuously-active (typically at
217 ~ 6 spikes s^{-1}), and some display rhythmic 'phasic' firing (69). Phasic firing is characterised by
218 bursts of activity that last more than 15 s, after which bursts stop randomly (70). Each burst is
219 followed by inactivity for at least 10 s, after which the next burst starts randomly (70). At burst
220 onset, vasopressin MNCs can reach firing rates of $\sim 15 - 25$ spikes s^{-1} for the first 5 – 10 s, before
221 spike frequency adaptation occurs to a steady-state firing rate of ~ 6 spikes s^{-1} for the remainder of
222 the burst (71, 72).

223 Of the different activity patterns recorded in vasopressin MNCs, phasic bursting is the most efficient
 224 pattern for vasopressin secretion into the circulation because vasopressin secretion is maximal at ~13
 225 spikes s⁻¹ (73, 74), which is typically only achieved by phasically firing MNCs and only during the
 226 first 5 – 10 s of each phasic burst. Vasopressin secretion from the posterior pituitary gland rapidly
 227 fatigues during continuous stimulation, but this fatigue is reversed when stimulation is stopped for a
 228 few tens of seconds (75). Hence, vasopressin MNCs firing continuously at high frequency do not
 229 secrete as much vasopressin into the circulation as do phasic MNCs firing at the same frequency
 230 because the silent periods between bursts in phasic MNCs reset the system for efficient vasopressin
 231 secretion at the onset of the next burst, when the typical firing frequency is again in the range that is
 232 most efficient for vasopressin secretion. The importance of phasic activity for efficient vasopressin
 233 secretion into the circulation is highlighted by the changes in activity patterning that occur under
 234 chronically stimulated conditions, such as prolonged osmotic stimulation. While burst duration does
 235 increase during shorter periods of stimulation, prolonged osmotic stimulation leads to an increase in
 236 firing rate within bursts while the burst duration and inter-burst interval remain similar to those seen
 237 under basal conditions (76-79).
 238 V1aR antagonists consistently increase the activity of phasic MNCs when administered into the
 239 SON (80), suggesting that somato-dendritic vasopressin mediates feedback inhibition of
 240 vasopressin MNCs via V1aR activation (Figure 5). This feedback inhibition likely involves direct
 241 autocrine actions on the MNC that secretes vasopressin because V1aR activation reduces excitatory
 242 postsynaptic potential amplitude in vasopressin MNCs (81). However, autocrine activation of
 243 V1aRs does not mediate autoregulation of vasopressin MNC activity alone because vasopressin
 244 also increases inhibitory postsynaptic potential frequency (82) via stimulation of astrocytic
 245 adenosine triphosphate (ATP) release, which acts as a gliotransmitter at P2X receptors on
 246 presynaptic GABA neurones to increase GABA release (67). Hence, somato-dendritic vasopressin
 247 secretion appears to contribute to the generation of phasic activity in vasopressin MNCs via a
 248 combination of autocrine actions on the MNC from which secretion occurs and paracrine actions
 249 on nearby cells that modulate the activity of the MNC from which secretion occurs.

250 While V1aR activation mediates autocrine and paracrine inhibition of phasic activity, local
 251 application of exogenous vasopressin was first reported to inhibit highly active phasic MNCs and
 252 stimulate weakly active phasic MNCs (83). Vasopressin MNCs also express V1bR (68), and while
 253 it has yet to be determined whether V1bR activation also contributes to autocrine regulation of
 254 vasopressin MNCs, it might underpin the excitatory effects of vasopressin evident in weakly active
 255 vasopressin MNCs. Regardless of the vasopressin excitation of weakly active phasic MNCs
 256 (perhaps via V1bR), such an action of endogenous vasopressin would presumably increase
 257 peripheral vasopressin secretion to cause robust vasoconstriction that is not present under basal
 258 conditions (84). Hence, it seems likely that any contribution of somato-dendritic vasopressin
 259 secretion to peripheral vasopressin secretion by feedback excitation is overridden by the V1aR-
 260 mediated feedback inhibition.

261 While, somato-dendritic vasopressin secretion functions as a negative feedback regulator of
 262 vasopressin MNC activity at the single cell level, the important output of the system is overall
 263 hormone secretion, which depends on the integrated activity of the MNCs at a population level
 264 (10). Some of the earliest work on somato-dendritic secretion showed that osmotic stimulation of
 265 vasopressin MNCs increases vasopressin levels in the circulation before levels increase in the SON
 266 (85), which is consistent with somato-dendritic vasopressin secretion acting as a negative feedback
 267 regulator of vasopressin MNC activity at a population level to modulate overall vasopressin
 268 secretion into the circulation.

269 **Autocrine/paracrine modulation of vasopressin magnocellular neurosecretory cell activity by** 270 **co-secreted transmitters**

271 Vasopressin MNCs also synthesise and secrete a number of other neurotransmitters and
 272 neuromodulators, including apelin (86), ATP (87), carbon monoxide (CO) (88), dynorphin (89),
 273 endocannabinoids (90-92), galanin (93), neuroendocrine regulatory peptides (NERPs) (94, 95),
 274 nitric oxide (NO) (96), PACAP (97) and secretin (13).

Most neuropeptides synthesised by MNCs are packaged within the same dense core vesicles as either vasopressin or oxytocin. However, apelin and galanin are differentially packaged in vasopressin MNCs. Apelin is packaged in dense core vesicles that do not contain vasopressin (98). While galanin is also packaged in some dense core vesicles that also contain vasopressin, it is also packaged in others that do not contain vasopressin and some dense core vesicles contain vasopressin but no galanin. Presumably, differential packaging in dense core vesicles might allow for secretion of separate pools that contain vasopressin or their co-expressed neuropeptides. Indeed, dense core vesicles containing galanin alone are trafficked to the dendrites while those that contain only vasopressin are trafficked to the axon terminals in the posterior pituitary gland (93).

Vasopressin MNCs express apelin receptors (APJ receptors) (99) and centrally-administered apelin inhibits vasopressin MNCs (86) to decrease basal vasopressin secretion (86, 100). However, systemic apelin administration increases vasopressin secretion (101) and chronic infusion of apelin into the PVN also increases vasopressin secretion (102). Furthermore, administration of apelin directly into the SON increases the activity of phasic MNCs (and presumably vasopressin secretion into the circulation) via non-specific cation channel activation, but reduces somato-dendritic vasopressin secretion (99), which presumably weakens vasopressin-mediated autoregulation to disinhibit and thus further excite vasopressin MNCs.

Vasopressin MNCs express galanin receptor-1 (Gal-R1) on their somata and dendrites (103). While centrally-administered galanin increases vasopressin secretion into the circulation *in vivo* (104, 105), it inhibits vasopressin secretion from isolated neurohypophyses or hypothalamo-neurohypophysial explants *in vitro* (106), suggesting that the direct effects of galanin are inhibitory, despite the reduced somato-dendritic vasopressin secretion. Indeed, galanin directly inhibits vasopressin MNCs *in vitro* by inducing hyperpolarization and reducing the slow afterdepolarisation (sADP) (107), which is a prominent excitatory post-spike potential in vasopressin MNCs (108). Galanin also reduces EPSC frequency (109), suggesting that it might also have paracrine effects after somato-dendritic secretion by retrograde inhibition of excitatory synaptic transmission.

Similarly to vasopressin receptors, κ -opioid receptors (KORs) are expressed in the membranes of vasopressin-containing dense core vesicles (110) and unlike apelin and galanin, the endogenous opioid peptide (EOP) ligand for KORs, dynorphin, is packaged with vasopressin in the same dense core vesicles (111). Hence, KORs newly-trafficked to the vasopressin MNC plasma membrane will be exposed to high concentrations of dynorphin upon somato-dendritic secretion of dense core vesicles. KOR agonists inhibit vasopressin MNCs *in vivo* (69, 112) and *in vitro* (113, 114). More importantly, antagonism of SON KORs increases burst duration in phasic MNCs under basal conditions *in vivo* (69, 70, 115) and *in vitro* (70, 116), showing that an endogenous KOR agonist inhibits phasic bursts. Phasic bursts are underpinned by the summation of sADPs to form a plateau potential that maintains a depolarised membrane potential to sustain further firing during bursts, and KOR activation causes activity-dependent sADP inhibition (116) to progressively decrease the plateau potential amplitude, which eventually leads to burst termination (70). Furthermore, KOR desensitisation prevents phasic activity in vasopressin MNCs, even when intensely stimulated (69). Hence, an endogenous KOR agonist inhibits phasic MNCs by autocrine inhibition of the sADP in the MNC from which dynorphin is secreted and this inhibition appears to be necessary for the expression of phasic activity by vasopressin MNCs.

In addition to sADP inhibition, KOR agonists reduce EPSP and IPSP amplitude (114, 117) and the delayed rectifier potassium current (118), while increasing the transient A-type potassium current (118) in vasopressin MNCs, although it has yet to be established whether these effects also contribute to the generation of phasic activity.

While KOR activation inhibits continuously-active vasopressin MNCs, KOR antagonism does not affect continuously-active vasopressin MNCs, even when they are strongly excited (76). Hence it appears that continuously-active vasopressin MNCs express KORs but do not release sufficient dynorphin to affect activity. Some vasopressin MNCs display irregular activity and these MNCs appear to be even more strongly excited by KOR antagonism than phasic MNCs (76). Taken together, this pattern-dependent sensitivity to KOR inhibition suggests that somato-dendritic dynorphin

secretion might determine the firing pattern of vasopressin MNCs and that transitions between firing patterns in individual vasopressin MNCs might result from changes in somato-dendritic dynorphin secretion (119).

MNCs also express receptors for pituitary adenylate cyclase-activating polypeptide (PACAP), which they also synthesise and secrete (97). PACAP increases somato-dendritic vasopressin secretion (120) by a direct depolarisation through activation of non-specific cation channels (121).

Neuroendocrine regulatory peptides (NERPs) 1 – 3 are packaged with vasopressin in the SON and PVN (94, 95). NERP-1 has paracrine effects on vasopressin MNC activity by retrograde inhibition of excitatory synaptic transmission whereas paracrine inhibition of vasopressin MNCs by NERP-2 is mediated by activation of upstream GABAergic interneurons that inhibit glutamatergic neurons that project to vasopressin MNCs (122). In contrast to NERPs 1 and 2, NERP-3 stimulates vasopressin secretion from the isolated posterior pituitary gland (95).

Vasopressin MNCs express secretin receptors and central secretin administration increases plasma vasopressin concentrations (13), suggesting that somato-dendritic secretin might stimulate systemic vasopressin secretion. However, secretin is also released from afferent inputs to the SON (123) and systemic secretin administration excites vasopressin (and oxytocin) MNCs via noradrenergic afferent inputs (124), suggesting that its actions might be mediated by afferent inputs rather than somato-dendritic secretion. While its role as a neurohypophysial hormone has not yet been definitively established, secretin is expressed in the posterior pituitary gland (13) and increases insertion of aquaporin-2 into the luminal membrane of the kidney to increase water reabsorption (125). Hence, secretin synthesised by vasopressin MNCs might act as a neurohypophysial hormone after secretion from the posterior pituitary gland rather than as an autoregulatory factor secreted from the somata and dendrites.

Vasopressin MNCs express P2X and P2Y receptors (126, 127) and injection of ATP into the SON induces antidiuresis (128). ATP is packaged in vasopressin dense core vesicles (87). ATP depolarises MNCs (129) and increases vasopressin secretion from hypothalamo-neurohypophysial explants (130).

ATP also increases glutamate and GABA release at synapses on MNCs (131). Hence, somato-dendritic ATP secretion might excite vasopressin MNCs by autocrine actions on the MNC from which it is secreted and paracrine actions on afferent inputs to the MNC from which it is secreted. However, MNCs are also excited by ATP released by astrocytes as a gliotransmitter (67, 132) as well as by ATP released from noradrenergic afferent inputs (133).

While somato-dendritic ATP secretion might modulate vasopressin MNC activity, ATP is rapidly catabolised to adenosine in the extracellular space (134). Vasopressin MNCs express adenosine A1 and A2A receptors (135) and A1 receptor antagonism excites phasic MNCs *in vivo*, but does not affect the firing rate of continuously-active vasopressin MNCs (136). A1 receptor antagonism reduces activity-dependent inhibition of EPSCs and IPSCs (137) as well as activity-dependent enhancement of the medium afterhyperpolarisation (mAHP) in vasopressin MNCs (138). mAHP activation induces spike frequency adaptation at the onset of phasic bursts (139) and so endogenous adenosine enhancement of the mAHP increases spike frequency adaptation, thereby shortening bursts in phasic MNCs (136). While A2 receptor activation depolarises MNCs to increase firing rate (135), vasopressin MNCs are inhibited when adenosine uptake is blocked. Hence, the overall effect of endogenous adenosine appears to be vasopressin MNC inhibition (140).

In addition to somato-dendritic exocytosis, vasopressin MNCs also release the gaseous transmitters, NO and carbon monoxide (CO), by diffusion after synthesis by NO synthase (NOS) and haem-oxygenase I, respectively. NO inhibits vasopressin MNCs (141, 142) by increasing IPSC amplitude and frequency (142, 143), whereas CO excites vasopressin MNCs (88).

Somato-dendritic modulation of vasopressin MNC activity appears to impact hormone secretion into the circulation through a complex interplay of excitatory (apelin, PACAP, ATP, NO and perhaps secretin) and inhibitory (galanin, dynorphin, NERPs, adenosine and CO) feedback that might fine tune the activity of individual MNCs to prevent any one MNC bearing too much of the secretory load for too long under basal conditions. It seems counter-intuitive that the autoregulatory effects of (at least some) co-secreted transmitters appear greater than that of vasopressin itself,

which is secreted in vastly greater quantities. Perhaps the autoregulatory effects of co-secreted transmitters are magnified by activation of both paracrine and autocrine mechanisms. In addition, it is possible that vasopressin's major role is paracrine inhibition of the population as a whole to prevent over-secretion of vasopressin into the circulation in response to perturbations of body fluid balance and/or blood pressure/volume.

Autocrine/paracrine modulation of oxytocin magnocellular neurosecretory cell activity

Similar to somato-dendritic vasopressin secretion, somato-dendritic oxytocin secretion also has autocrine and paracrine actions that modulate oxytocin MNC activity. In contrast to vasopressin, the autocrine and paracrine effects of oxytocin are arranged in series rather than in parallel; somato-dendritic oxytocin secretion activates oxytocin receptors (OTRs) on oxytocin MNCs to increase intracellular calcium, which has various actions including the release of endocannabinoids that cause retrograde inhibition of excitatory synaptic transmission under basal conditions (90). While the oxytocin-stimulated retrograde endocannabinoid suppression of excitatory synaptic input is expected to inhibit oxytocin MNCs (Figure 6), the best characterised effects of somato-dendritic oxytocin secretion is excitation, but only under specific (patho)physiological conditions. Hence, it has been proposed that endocannabinoid inhibition might occur over a longer timescale than autocrine effects of oxytocin to shape activity patterning in oxytocin MNCs during birth and lactation (144), but it has yet to be established whether this occurs in vivo. Alternatively, the excitatory effects of oxytocin might involve a switch from endocannabinoid inhibition to excitation during pregnancy, perhaps by enhanced expression/activation of excitatory transient receptor potential vanilloid-1 channels (145) for which the endocannabinoid, anandamide, is an endogenous ligand (146). Additionally, spillover of the endocannabinoid 2-arachidonoylglycerol from glutamate onto GABA synapses and a resulting suppression of inhibitory synaptic input has been observed in MNCs following glial retraction induced by salt loading (91). This endocannabinoid spillover could also occur with glial retraction during parturition and lactation to reduce inhibitory synaptic transmission, although this remains to be determined. Finally, it is also possible that oxytocin-stimulated endocannabinoid retrograde

modulation of excitatory and inhibitory synapses might be overridden by changes in the postsynaptic properties of oxytocin MNCs (147, 148), increased excitatory afferent inputs (149-153), or a switch in GABA signalling from inhibitory to excitatory, or less inhibitory, during pregnancy (64, 154).

Oxytocin MNCs typically exhibit continuous activity at $\sim 1 - 5$ spikes s^{-1} under basal conditions to maintain circulating oxytocin concentrations of $\sim 1 - 3$ pg ml^{-1} , with higher concentrations during sleep (2). However, oxytocin is best known for its stimulation of rhythmic uterine contraction during birth and of episodic milk ejection during suckling. Uterine contractions and mammary duct contraction each occur at intervals of several minutes and each contraction is triggered by a coordinated, high frequency burst of activity across the population of oxytocin MNCs (155) to secrete a discrete pulse of oxytocin into the circulation, which transiently increases intrauterine pressure during birth (156) and intramammary pressure during suckling (157).

Somato-dendritic oxytocin secretion increases immediately preceding each burst of activity in lactating rats (158) and bursts are blocked by OTR antagonist administration (159), as is the rise in somato-dendritic oxytocin secretion (160), suggesting that somato-dendritic oxytocin secretion is required for bursts to occur. However, the mechanisms by which somato-dendritic oxytocin secretion promotes burst firing in oxytocin MNCs are not fully understood and it is likely not the only contributor; synchronised volleys of EPSPs (161), rebound depolarization after bursts of IPSPs (162) and enhancement of the sADP (147, 163) have all been proposed to trigger or sustain firing during bursts in oxytocin MNCs.

In brain slices from male rats that do not normally fire bursts, α_1 -adrenoceptor activation in low calcium can induce bursts in oxytocin MNCs reminiscent of those seen during birth and milk ejection (164). Hence, noradrenergic inputs might trigger bursts. Consistent with this hypothesis, noradrenergic innervation of oxytocin MNCs is increased in late pregnancy (152) and these afferent inputs are activated during birth (55) to increase noradrenaline release into the SON (165).

1
2
3 429 Noradrenergic stimulation of bursts in oxytocin MNCs might be mediated by somato-dendritic
4
5 430 oxytocin release because noradrenergic receptor stimulation is required for suckling-induced somato-
6
7 431 dendritic oxytocin release (58), which might be part of a positive feedback loop that builds towards
8
9 432 bursts during continuous suckling. Indeed, burst-like activity can also be induced in virgin rats *in vivo*
10
11 433 by coordinated activation of neighbouring oxytocin MNCs, which induces priming of somato-
12
13 434 dendritic dense core vesicles for subsequent secretion (28). Once primed, high-frequency electrical
14
15 435 stimulation induces bursting in oxytocin MNCs of virgin rats (28). Hence, continuous suckling might
16
17 436 trigger tonic noradrenaline release onto oxytocin MNCs that triggers increasing somato-dendritic
18
19 437 oxytocin secretion, which could prime further somato-dendritic oxytocin secretion until a tipping-
20
21 438 point is reached to induce each burst.

22
23
24 439 In addition to facilitating burst firing in individual oxytocin MNCs, somato-dendritic oxytocin
25
26 440 secretion might also help coordinate the timing of bursts across the population of oxytocin MNCs.
27
28 441 Oxytocin injection into one SON increases the frequency of milk ejection bursts in the contralateral
29
30 442 SON (166). Oxytocin MNCs have 1 – 3 dendrites (148) that are normally separated from
31
32 443 neighbouring dendrites by astrocytic processes. However, in late pregnancy and lactation, astrocytes
33
34 444 withdraw their processes from between oxytocin MNC dendrites, which then form bundles of ~ 10
35
36 445 closely apposed dendrites (167, 168). A mathematical model in which oxytocin MNCs send each
37
38 446 dendrite to different dendritic bundles to form a sparse network of interactions emulates burst firing in
39
40 447 which each burst is initiated randomly at any of the dendritic bundles and spreads rapidly through the
41
42 448 oxytocin MNC population (144).

43
44
45 449 However, this model does not account for coordination of bursts across the bilateral SONs and PVNs
46
47 450 which might be mediated by noradrenergic inputs that project bilaterally to the SON (169). Indeed,
48
49 451 sectioning the optic chiasm or mammillary body disrupts co-ordination of bursts between oxytocin
50
51 452 MNCs in the left and right SON, suggesting that burst coordination across the magnocellular nuclei
52
53 453 involves projections through these areas (169, 170). Furthermore, the perinuclear zone that lies
54
55 454 immediately dorsal to the SON sends prominent projections to the SON (171) and PVN (172, 173)
56
57 455 that might also contribute to coordination of bursts across the four main magnocellular nuclei.
58
59
60

Autocrine/paracrine modulation of oxytocin magnocellular neurosecretory cell activity by co-secreted transmitters

Oxytocin MNCs also synthesise other transmitters that are likely secreted from their somata and dendrites, but the effects of these co-transmitters are not as well characterised as for those released from vasopressin MNCs.

Oxytocin MNCs express μ -opioid receptors (MORs) and KORs (174, 175), and MOR or KOR activation inhibits oxytocin MNCs (69, 176). While oxytocin MNCs synthesise μ - and κ -EOPs (177, 178), neither MOR nor KOR antagonists affect the activity of oxytocin MNCs *in vivo* (69, 179). Hence it appears that if EOPs undergo somato-dendritic secretion with oxytocin, they do not modulate oxytocin MNC activity to any appreciable extent under basal conditions. MOR-mediated EOP inhibition of somato-dendritic oxytocin secretion and oxytocin MNC activity is increased in late pregnancy (180), but this modulation is likely to be mediated by afferent inputs (181).

By contrast to the central actions of MOR activation, KOR activation appears to restrain secretion into the bloodstream at the posterior pituitary gland (182) and this effect also increases in late pregnancy (183). KOR restraint of oxytocin secretion might build up stores of oxytocin for birth and lactation and might potentiate secretion during bursts because KORs are desensitised on the day of birth (184, 185).

Oxytocin MNC dense core vesicles also contain ATP, which is presumably secreted along with oxytocin from the somata and dendrites. Co-secreted ATP does not modulate oxytocin MNC activity via adenosine receptor activation (136), it might excite oxytocin MNC via P2X receptor activation (130, 131).

Oxytocin MNCs also express NOS (96) and NO appears to restrain the activity of oxytocin MNCs, particularly under stimulated conditions (186, 187), suggesting that NO is an inhibitory autocrine/paracrine modulator of oxytocin MNC activity.

Remarkably, chronic MOR activation by the opioid alkaloid agonist, morphine, (but not EOPs (188)) induces tolerance and dependence in oxytocin MNCs (189). Tolerance is revealed as loss of inhibition to acute administration of morphine (176) and dependence is revealed by a sustained hyperexcitation upon withdrawal of chronic morphine administration (112). Somato-dendritic oxytocin secretion is increased during morphine withdrawal and OTR antagonism reduces morphine withdrawal-induced excitation of oxytocin MNCs (190). Hence, somato-dendritic oxytocin secretion appears to contribute to morphine withdrawal-induced excitation of oxytocin MNCs, although the mechanism by which it does so has yet to be identified.

Paracrine modulation of parvocellular paraventricular nucleus neurone activity by somato-dendritic secretion from magnocellular neurosecretory cells

While the SON essentially contains only MNCs (and glia and cells of the vasculature), the PVN also contains parvocellular neurones. Parvocellular neurones are sub-divided by their projections and functions: neurosecretory parvocellular neurones project to the hypothalamic median eminence, where they secrete hormones into the hypophysial portal blood vessels to control hormone secretion from the anterior pituitary gland; preautonomic parvocellular neurones project to the brainstem and spinal cord to modulate parasympathetic and sympathetic nervous system activity (191, 192); the remaining parvocellular neurones project to various brain areas to modulate behaviour.

It has long been hypothesised that somato-dendritic secretion from MNCs modulates the activity of parvocellular neurones but only recently has definitive evidence to support this hypothesis been generated for somato-dendritic vasopressin (38, 193, 194). For paracrine effects on other neuronal phenotypes to occur, first vasopressin (or oxytocin) must diffuse through the parenchyma to reach other neurones. The effective diffusion distance for somato-dendritic vasopressin was determined under basal conditions using Chinese hamster ovary cells transfected with human V1aRs and a calcium indicator to generate biosensor 'sniffer' cells with a threshold detection level of 0.5 nM and an EC_{50} of 7.2 nM for vasopressin (38). Using sniffer cells that were dispersed over the PVN in hypothalamic slices, it was shown that activation of an individual MNC induces sufficient somato-

1
2
3 506 dendritic vasopressin secretion to induce intracellular calcium increases for tens of seconds in sniffer
4
5 507 cells over 100 μm from the soma of the activated MNC (38). Similar results were seen using HEK-
6
7 508 293 sniffer cells in the SON (194). Hence, somato-dendritic vasopressin release from an individual
8
9 509 MNC diffuses through the PVN at sufficient concentration to activate V1aRs expressed on
10
11 510 parvocellular neurone somata or dendrites under basal conditions (38, 193). Remarkably, astrocytes
12
13 511 withdraw their processes from between MNCs when chronically stimulated (167, 168, 195, 196),
14
15 512 which reduces the tortuosity of the extracellular space and likely increases the effective diffusion
16
17 513 distance for somato-dendritic vasopressin and oxytocin through the parenchyma (197).
18
19
20
21 514 Some preautonomic parvocellular neurones project to the rostral ventrolateral medulla (RVLM) in the
22
23 515 brainstem, which projects to sympathetic ganglia to regulate sympathetic nerve activity (198).
24
25 516 RVLM-projecting parvocellular neurones express V1aRs and their dendrites are intermingled with
26
27 517 vasopressin MNC somata and dendrites in the PVN (193). Hence, the architecture is in place to
28
29 518 provide for somato-dendritic vasopressin modulation of autonomic function by paracrine modulation
30
31 519 of preautonomic neurones within the PVN (Figure 7). Activation of individual vasopressin MNCs by
32
33 520 uncaging NMDA increases action potential firing in RVLM-projecting parvocellular neurones beyond
34
35 521 100 μm from the activated MNC (193) and this activation is much more potent than that elicited by
36
37 522 action potential firing alone (38), suggesting that dendritic NMDARs might be a main driver of
38
39 523 somato-dendritic secretion. The responses of RVLM-projecting parvocellular neurones to vasopressin
40
41 524 MNC activation occur after a delay of several seconds, are blocked by superfusion of a V1aR
42
43 525 antagonist and are enhanced by a peptidase inhibitor (193). Most importantly, the depolarisation of
44
45 526 RVLM-projecting parvocellular neurones in response to vasopressin MNC activation is not affected
46
47 527 by blockade of action potential firing with tetrodotoxin. Taken together, the data demonstrate that the
48
49 528 excitation of RVLM-projecting parvocellular neurones is mediated by diffusion of somato-dendritic
50
51 529 vasopressin through the extracellular space of the PVN.
52
53
54
55
56 530 Given that activation of an individual vasopressin MNC can excite preautonomic neurones within a
57
58 531 radius of at least 100 μm , it appears likely that somato-dendritic secretion from the population of
59
60

MNCs, as occurs *in vivo*, could function as a population-to-population signal to recruit preautonomic neurones as a whole. Indeed, administration of a V1aR antagonist alone reduces preautonomic neurone activity and the more active the MNC, the more effective is the antagonist at reducing preautonomic neurone activity (193), suggesting that preautonomic neurones are under tonic modulation by vasopressin MNCs to coordinate the humoral (circulating vasopressin) and neuronal (sympathetic nerve activity) responses to changes in body fluid balance and to pathophysiological conditions, such as hypertension, myocardial infarction and chronic heart failure.

Hyperosmolality increases vasopressin secretion into the circulation by MNCs and increases renal sympathetic nerve activity, both of which increase water retention to protect body fluid balance. The mechanisms that underpin these responses to hyperosmolality are thoroughly reviewed elsewhere (199, 200). While these mechanisms occur in parallel, somato-dendritic vasopressin secretion likely coordinates the responses because intracarotid infusion of hyperosmotic saline causes a dose-dependent increase in renal sympathetic nerve activity that is accompanied by increased somato-dendritic vasopressin secretion, and bilateral injection of a V1aR antagonist into the PVN abolishes the renal sympathetic nerve activation (193). Hence, it appears that somato-dendritic vasopressin secretion coordinates the humoral and neuronal responses to increased osmolality.

PVN-driven sympathoexcitation is a key pathophysiological mechanism in hypertension (201, 202), acute myocardial infarction (192, 203) and heart failure (204, 205) that contributes to morbidity and mortality (206). Vasopressin MNCs are activated in hypertension (207, 208), acute myocardial infarction (209) and heart failure (210, 211). While the increased circulating vasopressin levels contribute directly to detrimental myocardial effects (212, 213), increased vasopressin MNC activity probably also increases somato-dendritic vasopressin secretion to contribute to the pathophysiological sympathoexcitation via activation of PVN preautonomic neurones.

While less well established than for vasopressin effects on preautonomic neurones, it appears that somato-dendritic oxytocin secretion might also modulate the activity of a neighbouring population of neurones within the PVN, corticotrophin-releasing hormone (CRH) neurones. Stress-induced CRH

secretion stimulates adrenocorticotrophic hormone secretion from the anterior pituitary gland (214) that, in turn, increases adrenal corticosteroid secretion to mediate the body's response to the stressor. CRH neurones express mRNA for OTRs (215) and oxytocin inhibits EPSC frequency, but not amplitude, in CRH neurones (216). Oxytocin MNC and CRH neurone dendrites are intermingled within the PVN (216), allowing for dendro-dendritic interactions between the two populations. Hence, somato-dendritic oxytocin might suppress CRH neurone excitability by presynaptic inhibition of excitatory synaptic inputs to reduce activation of the stress axis. However, OTR antagonism has no effect on CRH neurones in brain slices (216), suggesting that, unlike vasopressin modulation of preautonomic neurone activity, there is no OTR tone on CRH neurones, at least under *in vitro* basal conditions.

Paracrine modulation of arteriolar vasoconstriction in the supraoptic nucleus by somato-dendritic secretion from magnocellular neurosecretory cells

Classically, neuronal activity is thought to dilate arterioles and thereby increase local cerebral blood flow to meet the metabolic demands of active brain areas; this 'neurovascular coupling' is generally accepted to result from glutamatergic synaptic transmission that evokes release of vasoactive substances from neurones and astrocytes to relax vascular smooth muscle cells (217). However, vascular smooth muscle cells express V1aRs, providing a target for somato-dendritic vasopressin, at least within the SON and PVN. Vasoconstriction can be elicited in SON arterioles by stimulation of an individual vasopressin MNC, an effect that is blocked by V1aR antagonism (218) (Figure 7). Consistent with its effects on preautonomic neurones, somato-dendritic vasopressin can induce responses in arterioles beyond 100 μm from the activated MNC under basal conditions (218). Importantly, this V1aR-mediated vasoconstriction is over-ridden in hyperosmotic conditions by parallel release of NO, which causes vasodilation of local arterioles (218). Presumably the NO-induced vasodilation increases blood flow through the SON when increased vasopressin MNC activity is required to protect from further fluid loss and maintain blood pressure in the general circulation through vasopressin secretion from the posterior pituitary gland.

584 Paracrine modulation of neuronal activity beyond the paraventricular nucleus by somato-
585 dendritic secretion from magnocellular neurosecretory cells

586 While paracrine modulation of parvocellular neurones within the PVN by somato-dendritic secretion
 587 from MNCs is now well characterised, it has yet to be definitively established whether neuropeptides
 588 secreted from MNC somata and dendrites can affect the activity of neurones outside the PVN.

589 Nevertheless, there is some evidence that somato-dendritic oxytocin might act on neurones in brain
 590 areas relatively close to the SON and/or PVN, particularly for brain areas that receive little or no
 591 axonal projections from MNCs or from vasopressin or oxytocin parvocellular neurones.

592 The central effects of the primary anorexigenic hormone, leptin, are mediated by oxytocin, at least in
 593 part (219). Leptin is sensed by ARC POMC neurones that, as described above, project to the SON and
 594 PVN (61), where they secrete α -MSH to activate MC4-Rs (62) and thereby increase somato-dendritic
 595 oxytocin secretion (63, 220). Oxytocin inhibition of food intake is mediated, in part, by the
 596 ventromedial hypothalamus (VMH) because oxytocin injection into the VMH decreases food intake
 597 that is driven by energy balance rather than palatability (221). While OTRs are highly expressed in the
 598 VMH (222), there are essentially no oxytocin MNC axons in the VMH (219). Given that vasopressin
 599 released from a single MNC can activate cells over 100 μ m from the MNC soma from which it is
 600 secreted (38, 193) and the VMH is situated roughly between the SON and PVN, it is possible that
 601 somato-dendritic oxytocin release from MNCs could diffuse through the parenchyma in sufficient
 602 quantities to activate OTRs in the VMH, which have nanomolar affinity for oxytocin (223). It is also
 603 possible that OTR-expressing astrocytes in the SON and PVN could expand the spatial domain of the
 604 dendritically released oxytocin signalling by relaying the signals through astrocytic networks, as has
 605 been reported for vasopressin release from vasopressin MNC dendrites (67) and CRH neurone
 606 dendrites (224).

607 While oxytocin MNCs inhibit fear responses via axon collaterals to the CeA (17), it appears likely
 608 that SON somato-dendritic oxytocin secretion enhances social recognition via actions in the MeA.

609 The CeA contains oxytocin MNC axons collaterals but there are no oxytocin (MNC or parvocellular)

neurone axons in the MeA (17). OTRs are highly expressed in the MeA (222) and OTR antagonist injection into the MeA reduces social recognition induced by SON activation (225). However, MeA OTRs are not directly activated by oxytocin secreted into the CeA from MNC axon collaterals (17). The MeA lies immediately lateral to the SON and, even if somato-dendritic oxytocin secreted within the SON does not reach the MeA, some oxytocin MNC dendrites project to the MeA (225), which might deliver sufficient oxytocin to the MeA to promote social recognition. Alternatively, OTRs might be **activated** by vasopressin MNC axon collaterals in the MeA (19) because vasopressin has appreciable activity at OTRs (223).

Hormone-like modulation of neuronal activity in distant brain areas by somato-dendritic secretion from magnocellular neurosecretory cells

It has been hypothesised that somato-dendritic vasopressin and oxytocin from MNCs are a hormone-like signal in the brain with widespread effects on distant populations of neurones (226). However, accumulating evidence of the functional impact of MNC axon collaterals on behaviour via direct projections to distant brain areas (17-21) have led to this hypothesis being challenged (227).

The half-lives of vasopressin and oxytocin are ~20 min in the cerebrospinal fluid (CSF) (228), giving time for diffusion through the ventricular system, particularly downstream. However, vasopressin (at least) has a half-life of less than 1 min in the parenchyma (229). Given that the paracrine effects of somato-dendritic vasopressin on preautonomic neurones that are only ~100 μ m away is delayed by ~2 – 5 s (193), it appears unlikely that the neuropeptides could diffuse long distances through the brain to act as a hormone-like signal. However, dense core vesicle exocytosis is a slow process compared to microvesicle fusion at the synapse, with latencies of several seconds in hippocampal neurones (230), which could account for much of the latency of the preautonomic neurone response to somato-dendritic vasopressin secreted by MNCs in the PVN. Furthermore, vasopressin and oxytocin are secreted in sufficient quantities to be measured in dialysates collected from the SON and PVN (231), as well as in other brain areas (232-234) and in the cerebrospinal fluid (CSF) (235), suggesting that they diffuse through the parenchyma sufficiently to reach the CSF. Indeed, the microdialysis probes

used to measure vasopressin and oxytocin in many experiments have a recovery rate of <10% for vasopressin in the SON and PVN (236). Hence, it is likely that the actual concentrations of vasopressin and oxytocin present in the parenchyma and CSF are appreciably higher than those measured in dialysates. A further factor to be considered is retraction of astrocytic processes from around MNCs in dehydration and pregnancy (167, 168, 195, 196), which decreases tortuosity in the extracellular space, presumably allowing more ready escape of somato-dendritic vasopressin and oxytocin from the SON and PVN. It is difficult to imagine that the relatively sparse terminal fields of MNC axon collaterals and parvocellular vasopressin and oxytocin MNCs could release sufficient vasopressin and oxytocin to maintain the ambient levels of the neuropeptides found in the brain and CSF.

While there are clear examples of axon collaterals from sub-populations of MNCs affecting neuronal activity beyond the SON and PVN (237), this does not preclude the possibility that there is also long-distance inter-population signalling mediated by somato-dendritic volume transmission of vasopressin and oxytocin over a longer timescale, particularly in brain regions in which there are neuropeptide receptors but no neuropeptide axons, such as in the olfactory bulb. Nevertheless, there is still no compelling evidence for, or against, distal hormone-like signalling by somato-dendritic vasopressin and oxytocin transmission, although the levels of oxytocin and vasopressin present in the cerebrospinal fluid and in brain areas devoid of oxytocin and vasopressin axon terminals appear to be much higher than could be achieved from axon terminal release from MNC and parvocellular axon collaterals. The resolution of this debate remains an ongoing challenge for the field.

Concluding remarks

Autocrine/paracrine modulation of MNC activity by somato-dendritic vasopressin and oxytocin release has been extensively studied and is broadly accepted as a major function of somato-dendritic secretion from MNCs (238). While it is clear that co-secreted transmitters also modulate MNC activity, there is no evidence of paracrine actions on other neurones, even other MNCs. Indeed, somato-dendritic dynorphin terminates bursts in the MNC from which it is secreted (70), but there is

1
2
3 662 no correlation between burst termination in phasic MNCs in paired recordings using a single electrode
4
5 663 in which the recorded MNCs would be at most tens of micrometres apart (71), which is well within
6
7 664 the range of somato-dendritic vasopressin but evidently beyond the range of somato-dendritic
8
9 665 dynorphin. Hence, it is possible that co-secreted transmitters act as autocrine/paracrine modulators of
10
11 666 the individual MNC from which they are secreted, whereas the much higher levels of vasopressin or
12
13 667 oxytocin secreted modulate the activity of the population as a whole to regulate peripheral physiology.
14
15
16
17 668 Recently, compelling evidence has emerged that somato-dendritic secretion from MNCs modulates
18
19 669 arteriole diameter in the SON (218) as well as the activity of neurones, particularly RVLM-projecting
20
21 670 preautonomic neurones in the parvocellular PVN (193) (and perhaps also CRH neurones (216)). This
22
23 671 inter-population cross-talk between MNCs and preautonomic neurones likely coordinates the
24
25 672 hormonal (vasopressin secretion into the circulation) and neural (sympathetic nerve activation)
26
27 673 response to perturbations of body fluid balance and blood pressure/volume (239).
28
29
30 674 To date, there is no definitive evidence that somato-dendritic vasopressin and oxytocin have actions
31
32 675 beyond the SON and PVN and it remains to be determined whether these neuropeptides act as
33
34 676 hormone-like signals after secretion from MNC somata and dendrites. Nevertheless, while much of
35
36 677 the data are circumstantial, it appears likely that somato-dendritic oxytocin release modulates the
37
38 678 activity of neurones in some nearby brain areas that express OTRs but do not contain oxytocin (MNC
39
40 679 or parvocellular) neurone projections, specifically the VMH (219, 222) and MeA (222, 225). The
41
42 680 confirmation (or refutation) of the effects of somato-dendritic oxytocin on the VMH and/or MeA is
43
44 681 required to resolve this ongoing debate within the field.
45
46
47

48 **References**

49
50
51 683 1. Ludwig M. *Dendritic neurotransmitter release* New York: Kluwer Academic Publishers,
52
53 684 2004.
54
55 685 2. Brown CH. Magnocellular neurons and posterior pituitary function. *Compr Physiol.* 2016;
56
57 686 61701-41.
58
59
60

- 687 3. Verbalis JG, Mangione MP, Stricker EM. Oxytocin produces natriuresis in rats at
688 physiological plasma concentrations. *Endocrinology*. 1991; **128**(3): 1317-22.
- 689 4. Haanwinckel MA, Elias LK, Favaretto AL, Gutkowska J, McCann SM, Antunes-Rodrigues J.
690 Oxytocin mediates atrial natriuretic peptide release and natriuresis after volume expansion in the rat.
691 *Proc Natl Acad Sci U S A*. 1995; **92**(17): 7902-6.
- 692 5. Manaye KF, Lei DL, Tizabi Y, Davila-Garcia MI, Mouton PR, Kelly PH. Selective neuron
693 loss in the paraventricular nucleus of hypothalamus in patients suffering from major depression and
694 bipolar disorder. *J Neuropathol Exp Neurol*. 2005; **64**(3): 224-9.
- 695 6. Rhodes CH, Morrell JI, Pfaff DW. Immunohistochemical analysis of magnocellular elements
696 in rat hypothalamus: distribution and numbers of cells containing neurophysin, oxytocin, and
697 vasopressin. *J Comp Neurol*. 1981; **198**(1): 45-64.
- 698 7. Tweedle CD, Smithson KG, Hatton GI. Neurosecretory endings in the rat neurohypophysis
699 are en passant. *Exp Neurol*. 1989; **106**(1): 20-6.
- 700 8. Nordmann JJ, Morris JF. Method for quantitating the molecular content of a subcellular
701 organelle: hormone and neurophysin content of newly formed and aged neurosecretory granules.
702 *Proc Natl Acad Sci USA*. 1984; **81**(1): 180.
- 703 9. Leng G, Ludwig M. Neurotransmitters and peptides: whispered secrets and public
704 announcements. *J Physiol*. 2008; **586**(Pt 23): 5625-32.
- 705 10. Leng G, Brown C, Sabatier N, Scott V. Population dynamics in vasopressin cells.
706 *Neuroendocrinology*. 2008; **88**(3): 160-72.
- 707 11. Brown CH, Ruan M, Scott V, Tobin VA, Ludwig M. Multi-factorial somato-dendritic
708 regulation of phasic spike discharge in vasopressin neurons. *Prog Brain Res*. 2008; **170**: 219-28.
- 709 12. Meeker RB, Swanson DJ, Greenwood RS, Hayward JN. Ultrastructural distribution of
710 glutamate immunoreactivity within neurosecretory endings and pituicytes of the rat neurohypophysis.
711 *Brain Res*. 1991; **564**(2): 181-93.

13. Chu JY, Lee LTO, Lai CH, Vaudry H, Chan YS, Yung WH, Chow BK. Secretin as a neurohypophysial factor regulating body water homeostasis. *Proc Natl Acad Sci USA*. 2009; **106**(37): 15961-6.
14. Lemos JR, Ortiz-Miranda SI, Cuadra AE, Velazquez-Marrero C, Custer EE, Dad T, Dayanithi G. Modulation/physiology of calcium channel sub-types in neurosecretory terminals. *Cell Calcium*. 2012; **51**(3-4): 284-92.
15. Mason WT, Ho YW, Hatton GI. Axon collaterals of supraoptic neurones: anatomical and electrophysiological evidence for their existence in the lateral hypothalamus. *Neuroscience*. 1984; **11**(1): 169-82.
16. Hatton GI, Cobbett P, Salm AK. Extranuclear axon collaterals of paraventricular neurons in the rat hypothalamus: intracellular staining, immunocytochemistry and electrophysiology. *Brain Res Bull*. 1985; **14**(2): 123-32.
17. Knobloch HS, Charlet A, Hoffmann LC, Eliava M, Khrulev S, Cetin AH, Osten P, Schwarz MK, Seeburg PH, Stoop R, Grinevich V. Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron*. 2012; **73**(3): 553-66.
18. Menon R, Grund T, Zoicas I, Althammer F, Fiedler D, Biermeier V, Bosch OJ, Hiraoka Y, Nishimori K, Eliava M, Grinevich V, Neumann ID. Oxytocin signaling in the lateral septum prevents social fear during lactation. *Curr Biol*. 2018; **28**(7): 1066-78.e6.
19. Hernandez VS, Vazquez-Juarez E, Marquez MM, Jauregui-Huerta F, Barrio RA, Zhang L. Extra-neurohypophyseal axonal projections from individual vasopressin-containing magnocellular neurons in rat hypothalamus. *Front Neuroanat*. 2015; **9**:130.
20. Hernandez VS, Hernandez OR, Perez de la Mora M, Gomora MJ, Fuxe K, Eiden LE, Zhang L. Hypothalamic vasopressinergic projections innervate central amygdala GABAergic neurons: Implications for anxiety and stress coping. *Front Neural Circuits*. 2016; **10**:92.
21. Hernandez-Perez OR, Hernandez VS, Nava-Kopp AT, Barrio RA, Seifi M, Swinny JD, Eiden LE, Zhang L. A synaptically connected hypothalamic magnocellular vasopressin-locus coeruleus

- 738 neuronal circuit and its plasticity in response to emotional and physiological stress. *Front Neurosci.*
 739 2019; **13**196.
- 740 22. Yeo SH, Kyle V, Blouet C, Jones S, Colledge WH. Mapping neuronal inputs to Kiss1
 741 neurons in the arcuate nucleus of the mouse. *PLoS One.* 2019; **14**(3): e0213927.
- 742 23. Armstrong WE, Scholer J, McNeill TH. Immunocytochemical, Golgi and electron
 743 microscopic characterization of putative dendrites in the ventral glial lamina of the rat supraoptic
 744 nucleus. *Neuroscience.* 1982; **7**(3): 679-94.
- 745 24. Hatton GI. Emerging concepts of structure-function dynamics in adult brain: the
 746 hypothalamo-neurohypophysial system. *Prog Neurobiol.* 1990; **34**(6): 437-504.
- 747 25. Pow DV, Morris JF. Dendrites of hypothalamic magnocellular neurons release
 748 neurohypophysial peptides by exocytosis. *Neuroscience.* 1989; **32**(2): 435-9.
- 749 26. Ludwig M, Apps D, Menzies J, Patel JC, Rice ME. Dendritic release of neurotransmitters.
 750 *Compr Physiol.* 2017; **7**(1): 235-52.
- 751 27. Ludwig M, Bull PM, Tobin VA, Sabatier N, Landgraf R, Dayanithi G, Leng G. Regulation of
 752 activity-dependent dendritic vasopressin release from rat supraoptic neurones. *J Physiol.* 2005;
 753 **564**(2): 515-22.
- 754 28. Ludwig M, Sabatier N, Bull PM, Landgraf R, Dayanithi G, Leng G. Intracellular calcium
 755 stores regulate activity-dependent neuropeptide release from dendrites. *Nature.* 2002; **418**(6893): 85-
 756 9.
- 757 29. Dayanithi G, Forostyak O, Ueta Y, Verkhratsky A, Toescu EC. Segregation of calcium
 758 signalling mechanisms in magnocellular neurones and terminals. *Cell Calcium.* 2012; **51**(3-4): 293-9.
- 759 30. Fisher TE, Bourque CW. Calcium-channel subtypes in the somata and axon terminals of
 760 magnocellular neurosecretory cells. *Trends Neurosci.* 1996; **19**(10): 440-4.
- 761 31. Tobin VA, Douglas AJ, Leng G, Ludwig M. The involvement of voltage-operated calcium
 762 channels in somato-dendritic oxytocin release. *PLoS One.* 2011; **6**(10): e25366.

- 763 32. Bains JS, Ferguson AV. Activation of N-methyl-D-aspartate receptors evokes calcium spikes
764 in the dendrites of rat hypothalamic paraventricular nucleus neurons. *Neuroscience*. 1999; **90**(3): 885-
765 91.
- 766 33. de Kock CP, Wierda KD, Bosman LW, Min R, Koksma JJ, Mansvelder HD, Verhage M,
767 Brussaard AB. Somatodendritic secretion in oxytocin neurons is upregulated during the female
768 reproductive cycle. *J Neurosci*. 2003; **23**(7): 2726-34.
- 769 34. Brown CH, Scott V, Ludwig M, Leng G, Bourque CW. Somatodendritic dynorphin release:
770 orchestrating activity patterns of vasopressin neurons. *Biochem Soc Trans*. 2007; **35**(Pt 5): 1236-42.
- 771 35. Joe N, Scott V, Brown CH. Glial regulation of extrasynaptic NMDA receptor-mediated
772 excitation of supraoptic nucleus neurons during dehydration. *J Neuroendocrinol*. 2014; **26**: 35-42.
- 773 36. Sabatier N. alpha-Melanocyte-stimulating hormone and oxytocin: a peptide signalling
774 cascade in the hypothalamus. *J Neuroendocrinol*. 2006; **18**(9): 703-10.
- 775 37. Sabatier N, Caquineau C, Dayanithi G, Bull P, Douglas AJ, Guan XM, Jiang M, Van der PL,
776 Leng G. Alpha-melanocyte-stimulating hormone stimulates oxytocin release from the dendrites of
777 hypothalamic neurons while inhibiting oxytocin release from their terminals in the neurohypophysis. *J*
778 *Neurosci*. 2003; **23**(32): 10351-8.
- 779 38. Pitra S, Zhang M, Cauley E, Stern JE. NMDA receptors potentiate activity-dependent
780 dendritic release of neuropeptides from hypothalamic neurons. *J Physiol*. 2019; **597**(6): 1735-56.
- 781 39. Fleming TM, Scott V, Naskar K, Joe N, Brown CH, Stern JE. State-dependent changes in
782 astrocyte regulation of extrasynaptic NMDA receptor signalling in neurosecretory neurons. *J Physiol*.
783 2011; **589**(Pt 16): 3929-41.
- 784 40. Naskar K, Stern JE. A functional coupling between extrasynaptic NMDA receptors and A-
785 type K⁺ channels under astrocyte control regulates hypothalamic neurosecretory neuronal activity. *J*
786 *Physiol*. 2014; **592**: 2813-27.
- 787 41. Zhang M, Biancardi VC, Stern JE. An increased extrasynaptic NMDA tone inhibits A-type
788 K(+) current and increases excitability of hypothalamic neurosecretory neurons in hypertensive rats. *J*
789 *Physiol*. 2017; **595**(14): 4647-61.

- 790 42. Tobin VA, Hurst G, Norrie L, Dal Rio FP, Bull PM, Ludwig M. Thapsigargin-induced
791 mobilization of dendritic dense-cored vesicles in rat supraoptic neurons. *Eur J Neurosci.* 2004;
792 **19**(10): 2909-12.
- 793 43. Hurbin A, Boissin-Agasse L, Orcel H, Rabie A, Joux N, Desarmenien MG, Richard P, Moos
794 FC. The V_{1a} and V_{1b}, but not V₂, vasopressin receptor genes are expressed in the supraoptic nucleus of
795 the rat hypothalamus, and the transcripts are essentially colocalized in the vasopressinergic
796 magnocellular neurons. *Endocrinology.* 1998; **139**(11): 4701-7.
- 797 44. Meddle SL, Bishop VR, Gkoumassi E, van Leeuwen FW, Douglas AJ. Dynamic changes in
798 oxytocin receptor expression and activation at parturition in the rat brain. *Endocrinology.* 2007;
799 **148**(10): 5095-104.
- 800 45. Dayanithi G, Sabatier N, Widmer H. Intracellular calcium signalling in magnocellular
801 neurones of the rat supraoptic nucleus: understanding the autoregulatory mechanisms. *ExpPhysiol.*
802 2000; **85 Spec No75S**.
- 803 46. Zhang Z, Kindrat AN, Sharif-Naeini R, Bourque CW. Actin filaments mediate mechanical
804 gating during osmosensory transduction in rat supraoptic nucleus neurons. *J Neurosci.* 2007; **27**(15):
805 4008-13.
- 806 47. Prager-Khoutorsky M, Khoutorsky A, Bourque CW. Unique interweaved microtubule
807 scaffold mediates osmosensory transduction via physical interaction with TRPV1. *Neuron.* 2014;
808 **83**(4): 866-78.
- 809 48. Tobin VA, Ludwig M. The role of the actin cytoskeleton in oxytocin and vasopressin release
810 from rat supraoptic nucleus neurons. *J Physiol.* 2007; **582**(Pt 3): 1337-48.
- 811 49. Morris JF, Pow DV. Widespread release of peptides in the central nervous system:
812 quantitation of tannic acid-captured exocytoses. *AnatRec.* 1991; **231**(4): 437.
- 813 50. Zhou Q, Zhou P, Wang AL, Wu D, Zhao M, Sudhof TC, Brunger AT. The primed SNARE-
814 complexin-synaptotagmin complex for neuronal exocytosis. *Nature.* 2017; **548**(7668): 420-5.
- 815 51. Jurgutis P, Shuang R, Fletcher A, Stuenkel EL. Characterization and distribution of SNARE
816 proteins at neuroendocrine nerve endings. *Neuroendocrinology.* 1996; **64**(5): 379-92.

- 817 52. Pupier S, Leveque C, Marqueze B, Kataoka M, Takahashi M, Seagar MJ. Cysteine string
818 proteins associated with secretory granules of the rat neurohypophysis. *J Neurosci.* 1997; **17**(8): 2722-
819 7.
- 820 53. Miyata S, Takamatsu H, Maekawa S, Matsumoto N, Watanabe K, Kiyohara T, Hatton GI.
821 Plasticity of neurohypophysial terminals with increased hormonal release during dehydration:
822 ultrastructural and biochemical analyses. *J Comp Neurol.* 2001; **434**(4): 413.
- 823 54. Tobin V, Schwab Y, Lelos N, Onaka T, Pittman QJ, Ludwig M. Expression of exocytosis
824 proteins in rat supraoptic nucleus neurones. *J Neuroendocrinol.* 2012; **24**(4): 629-41.
- 825 55. Meddle SL, Leng G, Selvarajah JR, Bicknell RJ, Russell JA. Direct pathways to the
826 supraoptic nucleus from the brainstem and the main olfactory bulb are activated at parturition in the
827 rat. *Neuroscience.* 2000; **101**(4): 1013-21.
- 828 56. Brunton PJ, Russell JA. The expectant brain: adapting for motherhood. *Nat Rev Neurosci.*
829 2008; **9**(1): 11-25.
- 830 57. Lipschitz DL, Crowley WR, Bealer SL. Differential sensitivity of intranuclear and systemic
831 oxytocin release to central noradrenergic receptor stimulation during mid- and late gestation in rats.
832 *Am J Physiol Endocrinol Metab* 2004; **287**E523-8.
- 833 58. Bealer SL, Crowley WR. Noradrenergic control of central oxytocin release during lactation in
834 rats. *Am J Physiol.* 1998; **274**(3 Pt 1): E453-8.
- 835 59. Bealer SL, Armstrong WE, Crowley WR. Oxytocin release in magnocellular nuclei:
836 neurochemical mediators and functional significance during gestation. *Am J Physiol Regul Integr*
837 *Comp Physiol.* 2010; **299**(2): R452-28.
- 838 60. Onaka T, Ikeda K, Yamashita T, Honda K. Facilitative role of endogenous oxytocin in
839 noradrenaline release in the rat supraoptic nucleus. *Eur J Neurosci.* 2003; **18**(11): 3018-26.
- 840 61. Bagnol D, Lu XY, Kaelin CB, Day HE, Ollmann M, Gantz I, Akil H, Barsh GS, Watson SJ.
841 Anatomy of an endogenous antagonist: Relationship between Agouti-related protein and
842 proopiomelanocortin in brain. *J Neurosci.* 1999; **19**(18): Rc26.

62. Ladyman SR, Augustine RA, Scherf E, Phillipps HR, Brown CH, Grattan DR. Attenuated hypothalamic responses to alpha-melanocyte stimulating hormone during pregnancy in the rat. *J Physiol.* 2016; **594**(4): 1087-101.
63. Sabatier N, Leng G. Presynaptic actions of endocannabinoids mediate alpha-MSH-induced inhibition of oxytocin cells. *Am J Physiol Regul Integr Comp Physiol.* 2006; **290**(3): R577-84.
64. Augustine RA, Ladyman SR, Bouwer GT, Alyousif Y, Sapsford TJ, Scott V, Kokay IC, Grattan DR, Brown CH. Prolactin regulation of oxytocin neurone activity in pregnancy and lactation. *J Physiol.* 2017; **595**:591–605.
65. Sakata I, Nakano Y, Osborne-Lawrence S, Rovinsky SA, Lee CE, Perello M, Anderson JG, Coppari R, Xiao G, Lowell BB, Elmquist JK, Zigman JM. Characterization of a novel ghrelin cell reporter mouse. *Regul Pept.* 2009; **155**(1-3): 91-8.
66. Ishizaki S, Murase T, Sugimura Y, Kakiya S, Yokoi H, Tachikawa K, Arima H, Miura Y, Oiso Y. Role of ghrelin in the regulation of vasopressin release in conscious rats. *Endocrinology.* 2002; **143**(5): 1589-93.
67. Haam J, Halmos KC, Di S, Tasker JG. Nutritional state-dependent ghrelin activation of vasopressin neurons via retrograde trans-neuronal-glial stimulation of excitatory GABA circuits. *J Neurosci.* 2014; **34**(18): 6201-13.
68. Hurbin A, Orcel H, Alonso G, Moos F, Rabie A. The vasopressin receptors colocalize with vasopressin in the magnocellular neurons of the rat supraoptic nucleus and are modulated by water balance. *Endocrinology.* 2002; **143**(2): 456-66.
69. Brown CH, Ludwig M, Leng G. Kappa-opioid regulation of neuronal activity in the rat supraoptic nucleus *in vivo*. *J Neurosci.* 1998; **18**(22): 9480-8.
70. Brown CH, Leng G, Ludwig M, Bourque CW. Endogenous activation of supraoptic nucleus kappa-opioid receptors terminates spontaneous phasic bursts in rat magnocellular neurosecretory cells. *J Neurophysiol.* 2006; **95**(5): 3235-44.
71. Brown CH. Rhythmogenesis in vasopressin cells. *J Neuroendocrinol.* 2004; **16**(9): 727-39.

- 869 72. Brown CH, Bourque CW. Mechanisms of rhythmogenesis: insights from hypothalamic
870 vasopressin neurons. *Trends Neurosci.* 2006; **29**(2): 108-15.
- 871 73. Bicknell RJ. Optimizing release from peptide hormone secretory nerve terminals. *J Exp Biol.*
872 1988; **139**51-65.
- 873 74. Dutton A, Dyball RE. Phasic firing enhances vasopressin release from the rat
874 neurohypophysis. *J Physiol.* 1979; **290**(2): 433-40.
- 875 75. Bicknell RJ, Brown D, Chapman C, Hancock PD, Leng G. Reversible fatigue of stimulus-
876 secretion coupling in the rat neurohypophysis. *J Physiol.* 1984; **348**601-13.
- 877 76. Scott V, Bishop VR, Leng G, Brown CH. Dehydration-induced modulation of kappa-opioid
878 inhibition of vasopressin neurone activity. *J Physiol.* 2009; **587**5679-89.
- 879 77. Wakerley JB, Poulain DA, Brown D. Comparison of firing patterns in oxytocin- and
880 vasopressin-releasing neurones during progressive dehydration. *Brain Res.* 1978; **148**(2): 425-40.
- 881 78. Dyball RE, Pountney PS. Discharge patterns of supraoptic and paraventricular neurones in
882 rats given a 2 per cent NaCl solution instead of drinking water. *J Endocrinol.* 1973; **56**(1): 91-8.
- 883 79. Walters JK, Hatton GI. Supraoptic neuronal activity in rats during five days of water
884 deprivation. *Physiol Behav.* 1974; **13**(5): 661-7.
- 885 80. Ludwig M, Leng G. Autoinhibition of supraoptic nucleus vasopressin neurons *in vivo*: a
886 combined retrodialysis/electrophysiological study in rats. *Eur J Neurosci.* 1997; **9**(12): 2532-40.
- 887 81. Kombian SB, Mougnot D, Hirasawa M, Pittman QJ. Vasopressin preferentially depresses
888 excitatory over inhibitory synaptic transmission in the rat supraoptic nucleus *in vitro*. *J*
889 *Neuroendocrinol.* 2000; **12**(4): 361-7.
- 890 82. Hermes ML, Ruijter JM, Klop A, Buijs RM, Renaud LP. Vasopressin increases GABAergic
891 inhibition of rat hypothalamic paraventricular nucleus neurons *in vitro*. *J Neurophysiol.* 2000; **83**(2):
892 705-11.
- 893 83. Gouzenes L, Desarmenien MG, Hussy N, Richard P, Moos FC. Vasopressin regularizes the
894 phasic firing pattern of rat hypothalamic magnocellular vasopressin neurons. *J Neurosci.* 1998; **18**(5):
895 1879-85.

- 896 84. Yosten GL, Samson WK. Pressor doses of vasopressin result in only transient elevations in
897 plasma peptide levels. *Peptides*. 2012; **33**(2): 342-5.
- 898 85. Ludwig M, Callahan MF, Neumann I, Landgraf R, Morris M. Systemic osmotic stimulation
899 increases vasopressin and oxytocin release within the supraoptic nucleus. *J Neuroendocrinol*. 1994;
900 **6**(4): 369-73.
- 901 86. De Mota N, Reaux-Le Goazigo A, El Messari S, Chartrel N, Roesch D, Dujardin C, Kordon
902 C, Vaudry H, Moos F, Llorens-Cortes C. Apelin, a potent diuretic neuropeptide counteracting
903 vasopressin actions through inhibition of vasopressin neuron activity and vasopressin release. *Proc*
904 *Natl Acad Sci U S A*. 2004; **101**(28): 10464-9.
- 905 87. Poisner AM, Douglas WW. Adenosine triphosphate and adenosine triphosphatase in
906 hormone-containing granules of posterior pituitary gland. *Science*. 1968; **160**(824): 203-4.
- 907 88. Reis WL, Biancardi VC, Son S, Antunes-Rodrigues J, Stern JE. Enhanced expression of heme
908 oxygenase-1 and carbon monoxide excitatory effects in oxytocin and vasopressin neurones during
909 water deprivation. *J Neuroendocrinol*. 2012; **24**(4): 653-63.
- 910 89. Watson SJ, Akil H, Fischli W, Goldstein A, Zimmerman E, Nilaver G, Wimersma Griedanus
911 TB. Dynorphin and vasopressin: common localization in magnocellular neurons. *Science*. 1982;
912 **216**(4541): 85-7.
- 913 90. Hirasawa M, Schwab Y, Natah S, Hillard CJ, Mackie K, Sharkey KA, Pittman QJ.
914 Dendritically released transmitters cooperate via autocrine and retrograde actions to inhibit afferent
915 excitation in rat brain. *J Physiol*. 2004; **559**(Pt 2): 611-24.
- 916 91. Di S, Popescu IR, Tasker JG. Glial control of endocannabinoid heterosynaptic modulation in
917 hypothalamic magnocellular neuroendocrine cells. *J Neurosci*. 2013; **33**(46): 18331-42.
- 918 92. Iremonger KJ, Kuzmiski JB, Baimoukhametova DV, Bains JS. Dual regulation of anterograde
919 and retrograde transmission by endocannabinoids. *J Neurosci*. 2011; **31**(33): 12011-20.
- 920 93. Landry M, Vila-Porcile E, Hokfelt T, Calas A. Differential routing of coexisting
921 neuropeptides in vasopressin neurons. *Eur J Neurosci*. 2003; **17**(3): 579-89.

- 922 94. Yamaguchi H, Sasaki K, Satomi Y, Shimbara T, Kageyama H, Mondal MS, Toshinai K, Date
923 Y, González LJ, Shioda S, Takao T, Nakazato M, Minamino N. Peptidomic identification and
924 biological validation of neuroendocrine regulatory peptide-1 and -2. *J Biol Chem.* 2007; **282**(36):
925 26354-60.
- 926 95. Fujihara H, Sasaki K, Mishiro-Sato E, Ohbuchi T, Dayanithi G, Yamasaki M, Ueta Y,
927 Minamino N. Molecular characterization and biological function of neuroendocrine regulatory
928 peptide-3 in the rat. *Endocrinology.* 2012; **153**(3): 1377-86.
- 929 96. Kadowaki K, Kishimoto J, Leng G, Emson PC. Up-regulation of nitric oxide synthase (NOS)
930 gene expression together with NOS activity in the rat hypothalamo-hypophyseal system after chronic
931 salt loading: evidence of a neuromodulatory role of nitric oxide in arginine vasopressin and oxytocin
932 secretion. *Endocrinology.* 1994; **134**(3): 1011-7.
- 933 97. Gillard ER, Leon-Olea M, Mucio-Ramirez S, Coburn CG, Sanchez-Islas E, de Leon A,
934 Mussenden H, Bauce LG, Pittman QJ, Curras-Collazo MC. A novel role for endogenous pituitary
935 adenylate cyclase activating polypeptide in the magnocellular neuroendocrine system. *Endocrinology.*
936 2006; **147**(2): 791-803.
- 937 98. Reaux-Le Goazigo A, Morinville A, Burlet A, Llorens-Cortes C, Beaudet A. Dehydration-
938 induced cross-regulation of apelin and vasopressin immunoreactivity levels in magnocellular
939 hypothalamic neurons. *Endocrinology.* 2004; **145**(9): 4392-400.
- 940 99. Tobin VA, Bull PM, Arunachalam S, O'Carroll AM, Ueta Y, Ludwig M. The effects of apelin
941 on the electrical activity of hypothalamic magnocellular vasopressin and oxytocin neurons and
942 somatodendritic peptide release. *Endocrinology.* 2008; **149**(12): 6136-45.
- 943 100. Reaux A, De Mota N, Skultetyova I, Lenkei Z, El Messari S, Gallatz K, Corvol P, Palkovits
944 M, Llorens-Cortes C. Physiological role of a novel neuropeptide, apelin, and its receptor in the rat
945 brain. *J Neurochem.* 2001; **77**(4): 1085-96.
- 946 101. Hus-Citharel A, Bodineau L, Frugiere A, Joubert F, Bouby N, Llorens-Cortes C. Apelin
947 counteracts vasopressin-induced water reabsorption via cross talk between apelin and vasopressin
948 receptor signaling pathways in the rat collecting duct. *Endocrinology.* 2014; **155**(11): 4483-93.

102. Zhang F, Sun HJ, Xiong XQ, Chen Q, Li YH, Kang YM, Wang JJ, Gao XY, Zhu GQ. Apelin-13 and APJ in paraventricular nucleus contribute to hypertension via sympathetic activation and vasopressin release in spontaneously hypertensive rats. *Acta Physiol (Oxf)*. 2014; **212**(1): 17-27.
103. Burazin TC, Larm JA, Gundlach AL. Regulation by osmotic stimuli of galanin-R1 receptor expression in magnocellular neurones of the paraventricular and supraoptic nuclei of the rat. *J Neuroendocrinol*. 2001; **13**(4): 358-70.
104. Ciosek J, Cisowska A. Centrally administered galanin modifies vasopressin and oxytocin release from the hypothalamo-neurohypophyseal system of euhydrated and dehydrated rats. *J Physiol Pharmacol*. 2003; **54**(4): 625-41.
105. Cisowska-Maciejewska A, Ciosek J. Galanin influences vasopressin and oxytocin release from the hypothalamo-neurohypophyseal system of salt loaded rats. *J Physiol Pharmacol*. 2005; **56**(4): 673-88.
106. Wodowska J, Ciosek J. Galanin and galanin-like peptide modulate vasopressin and oxytocin release in vitro: the role of galanin receptors. *Neuropeptides*. 2014; **48**(6): 387-97.
107. Papas S, Bourque CW. Galanin inhibits continuous and phasic firing in rat hypothalamic magnocellular neurosecretory cells. *J Neurosci*. 1997; **17**(16): 6048-56.
108. Andrew RD, Dudek FE. Burst discharge in mammalian neuroendocrine cells involves an intrinsic regenerative mechanism. *Science*. 1983; **221**(4615): 1050-2.
109. Kozoriz MG, Kuzmiski JB, Hirasawa M, Pittman QJ. Galanin modulates neuronal and synaptic properties in the rat supraoptic nucleus in a use and state dependent manner. *J Neurophysiol*. 2006; **96**(1): 154-64.
110. Shuster SJ, Riedl M, Li X, Vulchanova L, Elde R. The kappa opioid receptor and dynorphin co-localize in vasopressin magnocellular neurosecretory neurons in guinea-pig hypothalamus. *Neuroscience*. 2000; **96**(2): 373-83.
111. Whitnall MH, Gainer H, Cox BM, Molineaux CJ. Dynorphin-A-(1-8) is contained within vasopressin neurosecretory vesicles in rat pituitary. *Science*. 1983; **222**(4628): 1137-9.

- 1
- 2
- 3 975 112. Ludwig M, Brown CH, Russell JA, Leng G. Local opioid inhibition and morphine
- 4
- 5 976 dependence of supraoptic nucleus oxytocin neurones in the rat *in vivo* *J Physiol*. 1997; **505**(Pt 1): 145-
- 6
- 7 977 52.
- 8
- 9 978 113. Inenaga K, Imura H, Yanaihara N, Yamashita H. Kappa-selective opioid receptor agonists
- 10
- 11 979 leumorphin and dynorphin inhibit supraoptic neurons in rat hypothalamic slice preparations. *J*
- 12
- 13 980 *Neuroendocrinol*. 1990; **23**89.
- 14
- 15 981 114. Inenaga K, Nagatomo T, Nakao K, Yanaihara N, Yamashita H. Kappa-selective agonists
- 16
- 17 982 decrease postsynaptic potentials and calcium components of action potentials in the supraoptic
- 18
- 19 983 nucleus of rat hypothalamus *in vitro*. *Neuroscience*. 1994; **58**(2): 331-40.
- 20
- 21
- 22 984 115. Brown CH, Ludwig M, Leng G. Temporal dissociation of the feedback effects of
- 23
- 24 985 dendritically co-released peptides on rhythmogenesis in vasopressin cells. *Neuroscience*. 2004;
- 25
- 26 986 **124**(1): 105-11.
- 27
- 28 987 116. Brown CH, Bourque CW. Autocrine feedback inhibition of plateau potentials terminates
- 29
- 30 988 phasic bursts in magnocellular neurosecretory cells of the rat supraoptic nucleus. *J Physiol*. 2004;
- 31
- 32 989 **557**949-60.
- 33
- 34 990 117. Iremonger KJ, Bains JS. Retrograde opioid signaling regulates glutamatergic transmission in
- 35
- 36 991 the hypothalamus. *J Neurosci*. 2009; **29**(22): 7349-58.
- 37
- 38
- 39 992 118. Muller W, Hallermann S, Swandulla D. Opioidergic modulation of voltage-activated K +
- 40
- 41 993 currents in magnocellular neurons of the supraoptic nucleus in rat. *J Neurophysiol*. 1999; **81**(4): 1617.
- 42
- 43 994 119. Scott V, Brown CH. State-dependent plasticity in vasopressin neurones: dehydration-induced
- 44
- 45 995 changes in activity patterning. *J Neuroendocrinol*. 2010; **22**(5): 343-54.
- 46
- 47 996 120. Shibuya I, Noguchi J, Tanaka K, Harayama N, Inoue U, Kabashima N, Ueta Y, Hattori Y,
- 48
- 49 997 Yamashita H. PACAP increases the cytosolic Ca²⁺ concentration and stimulates somatodendritic
- 50
- 51 998 vasopressin release in rat supraoptic neurons. *J Neuroendocrinol*. 1998; **10**(1): 31-42.
- 52
- 53 999 121. Shibuya I, Kabashima N, Tanaka K, Setiadji VS, Noguchi J, Harayama N, Ueta Y, Yamashita
- 54
- 55 1000 H. Patch-clamp analysis of the mechanism of PACAP-induced excitation in rat supraoptic neurones. *J*
- 56
- 57 1001 *Neuroendocrinol*. 1998; **10**(10): 759-68.
- 58
- 59
- 60

- 1002 122. Toshinai K, Saito T, Yamaguchi H, Sasaki K, Tsuchimochi W, Minamino N, Ueta Y,
1003 Nakazato M. Neuroendocrine regulatory peptide-1 and -2 (NERPs) inhibit the excitability of
1004 magnocellular neurosecretory cells in the hypothalamus. *Brain Res.* 2014; **1563**:52-60.
- 1005 123. Lee VH, Lee LT, Chu JY, Lam IP, Siu FK, Vaudry H, Chow BK. An indispensable role of
1006 secretin in mediating the osmoregulatory functions of angiotensin II. *FASEB J.* 2010; **24**(12): 5024-
1007 32.
- 1008 124. Velmurugan S, Brunton PJ, Leng G, Russell JA. Circulating secretin activates supraoptic
1009 nucleus oxytocin and vasopressin neurons via noradrenergic pathways in the rat. *Endocrinology.*
1010 2010; **151**(6): 2681-8.
- 1011 125. Chu JY, Chung SC, Lam AK, Tam S, Chung SK, Chow BK. Phenotypes developed in
1012 secretin receptor-null mice indicated a role for secretin in regulating renal water reabsorption. *Mol*
1013 *Cell Biol.* 2007; **27**(7): 2499-511.
- 1014 126. Xiang Z, Bo X, Oglesby I, Ford A, Burnstock G. Localization of ATP-gated P2X2 receptor
1015 immunoreactivity in the rat hypothalamus. *Brain Res.* 1998; **813**(2): 390-7.
- 1016 127. Song Z, Vijayaraghavan S, Sladek CD. ATP increases intracellular calcium in supraoptic
1017 neurons by activation of both P2X and P2Y purinergic receptors. *Am J Physiol Regul Integr Comp*
1018 *Physiol.* 2007; **292**(1): R423-31.
- 1019 128. Mori M, Tsushima H, Matsuda T. Antidiuretic effects of ATP induced by microinjection into
1020 the hypothalamic supraoptic nucleus in water-loaded and ethanol-anesthetized rats. *Jpn J Pharmacol.*
1021 1994; **66**(4): 445-50.
- 1022 129. Hiruma H, Bourque CW. P2 purinoceptor-mediated depolarization of rat supraoptic
1023 neurosecretory cells in vitro. *J Physiol.* 1995; **489** (Pt 3):805-11.
- 1024 130. Song Z, Gomes DA, Stevens W, Sladek CD. Multiple alpha1-adrenergic receptor subtypes
1025 support synergistic stimulation of vasopressin and oxytocin release by ATP and phenylephrine. *Am J*
1026 *Physiol Regul Integr Comp Physiol.* 2010; **299**(6): R1529-37.

- 1027 131. Vavra V, Bhattacharya A, Zemkova H. Facilitation of glutamate and GABA release by P2X
- 1028 receptor activation in supraoptic neurons from freshly isolated rat brain slices. *Neuroscience*. 2011;
- 1029 **188**1-12.
- 1030 132. Gordon GR, Baimoukhametova DV, Hewitt SA, Rajapaksha WR, Fisher TE, Bains JS.
- 1031 Norepinephrine triggers release of glial ATP to increase postsynaptic efficacy. *Nat Neurosci*. 2005;
- 1032 **8**(8): 1078-86.
- 1033 133. Day TA, Sibbald JR, Khanna S. ATP mediates an excitatory noradrenergic neuron input to
- 1034 supraoptic vasopressin cells. *Brain Res*. 1993; **607**(1-2): 341-4.
- 1035 134. Song Z, Sladek CD. Does conversion of ATP to adenosine terminate ATP-stimulated
- 1036 vasopressin release from hypothalamo-neurohypophyseal explants? *Brain Res*. 2005; **1047**(1): 105-
- 1037 11.
- 1038 135. Ponzio TA, Wang YF, Hatton GI. Activation of adenosine A2A receptors alters postsynaptic
- 1039 currents and depolarizes neurons of the supraoptic nucleus. *Am J Physiol Regul Integr Comp Physiol*.
- 1040 2006; **291**(2): R359-66.
- 1041 136. Bull PM, Brown CH, Russell JA, Ludwig M. Activity-dependent feedback modulation of
- 1042 spike patterning of supraoptic nucleus neurons by endogenous adenosine. *Am J Physiol Regul Integr*
- 1043 *Comp Physiol*. 2006; **291**(1): R83-90.
- 1044 137. Oliet SH, Poulain DA. Adenosine-induced presynaptic inhibition of IPSCs and EPSCs in rat
- 1045 hypothalamic supraoptic nucleus neurones. *J Physiol*. 1999; **520**(Pt 3): 815-25.
- 1046 138. Ruan M, Brown CH. Feedback inhibition of action potential discharge by endogenous
- 1047 adenosine enhancement of the medium afterhyperpolarization. *J Physiol*. 2009; **587**(Pt 5): 1043-56.
- 1048 139. Kirkpatrick K, Bourque CW. Activity dependence and functional role of the apamin-sensitive
- 1049 K⁺ current in rat supraoptic neurones *in vitro*. *J Physiol*. 1996; **494**(Pt 2): 389-98.
- 1050 140. Ponzio TA, Hatton GI. Adenosine postsynaptically modulates supraoptic neuronal
- 1051 excitability. *J Neurophysiol*. 2005; **93**(1): 535-47.
- 1052 141. Liu QS, Jia YS, Ju G. Nitric oxide inhibits neuronal activity in the supraoptic nucleus of the
- 1053 rat hypothalamic slices. *Brain Res Bull*. 1997; **43**(2): 121-5.

- 1054 142. Stern JE, Ludwig M. NO inhibits supraoptic oxytocin and vasopressin neurons via activation
1055 of GABAergic synaptic inputs. *Am J Physiol.* 2001; **280**(6): R1815-22.
- 1056 143. Ozaki M, Shibuya I, Kabashima N, Isse T, Noguchi J, Ueta Y, Inoue Y, Shigematsu A,
1057 Yamashita H. Preferential potentiation by nitric oxide of spontaneous inhibitory postsynaptic currents
1058 in rat supraoptic neurones. *J Neuroendocrinol.* 2000; **12**(3): 273-81.
- 1059 144. Rossoni E, Feng J, Tirozzi B, Brown D, Leng G, Moos F. Emergent synchronous bursting of
1060 oxytocin neuronal network. *PLoS Comput Biol.* 2008; **4**(7): e1000123.
- 1061 145. Sharif Naeini R, Witty MF, Seguela P, Bourque CW. An N-terminal variant of Trpv1 channel
1062 is required for osmosensory transduction. *Nat Neurosci.* 2006; **9**(1): 93-8.
- 1063 146. Zygmunt PM, Petersson J, Andersson DA, Chuang H, Sorgard M, Di Marzo V, Julius D,
1064 Hogestatt ED. Vanilloid receptors on sensory nerves mediate the vasodilator action of anandamide.
1065 *Nature.* 1999; **400**(6743): 452-7.
- 1066 147. Stern JE, Armstrong WE. Changes in the electrical properties of supraoptic nucleus oxytocin
1067 and vasopressin neurons during lactation. *J Neurosci.* 1996; **16**(16): 4861-71.
- 1068 148. Stern JE, Armstrong WE. Reorganization of the dendritic trees of oxytocin and vasopressin
1069 neurons of the rat supraoptic nucleus during lactation. *J Neurosci.* 1998; **18**(3): 841-53.
- 1070 149. El Majdoubi M, Poulain DA, Theodosia DT. The glutamatergic innervation of oxytocin- and
1071 vasopressin-secreting neurons in the rat supraoptic nucleus and its contribution to lactation-induced
1072 synaptic plasticity. *Eur J Neurosci.* 1996; **8**(7): 1377.
- 1073 150. Gies U, Theodosia DT. Synaptic plasticity in the rat supraoptic nucleus during lactation
1074 involves GABA innervation and oxytocin neurons: a quantitative immunocytochemical analysis.
1075 *Journal of Neuroscience.* 1994; **14**(5 Pt 1): 2861.
- 1076 151. Stern JE, Hestrin S, Armstrong WE. Enhanced neurotransmitter release at glutamatergic
1077 synapses on oxytocin neurones during lactation in the rat. *J Physiol.* 2000; **526**(Pt 1): 109-14.
- 1078 152. Michaloudi HC, El Majdoubi M, Poulain DA, Papadopoulos GC, Theodosia DT. The
1079 noradrenergic innervation of identified hypothalamic magnocellular somata and its contribution to
1080 lactation-induced synaptic plasticity. *J Neuroendocrinol.* 1997; **9**(1): 17-23.

1
2
3 1081 153. Seymour AJ, Scott V, Augustine RA, Bouwer GT, Campbell RE, Brown CH. Development of
4
5 1082 an excitatory kisspeptin projection to the oxytocin system in late pregnancy. *J Physiol.* 2017; **595**(3):
6
7 1083 825-38.
8
9 1084 154. Lee SW, Kim YB, Kim JS, Kim WB, Kim YS, Han HC, Colwell CS, Cho YW, Kim YI.
10
11 1085 GABAergic inhibition is weakened or converted into excitation in the oxytocin and vasopressin
12
13 1086 neurons of the lactating rat. *Mol Brain.* 2015; **8**34.
14
15 1087 155. Belin V, Moos F, Richard P. Synchronization of oxytocin cells in the hypothalamic
16
17 1088 paraventricular and supraoptic nuclei in suckled rats: direct proof with paired extracellular recordings.
18
19 1089 *Exp Brain Res.* 1984; **57**(1): 201-3.
20
21
22 1090 156. Douglas A, Scullion S, Antonijevic I, Brown D, Russell J, Leng G. Uterine contractile
23
24 1091 activity stimulates supraoptic neurons in term pregnant rats via a noradrenergic pathway.
25
26 1092 *Endocrinology.* 2001; **142**(2): 633-44.
27
28 1093 157. Augustine RA, Seymour AJ, Campbell RE, Grattan DR, Brown CH. Integrative neuro-
29
30 1094 humoral regulation of oxytocin neuron activity in pregnancy and lactation. *J Neuroendocrinol.* 2018;
31
32 1095 **30**e12569.
33
34 1096 158. Moos F, Poulain DA, Rodriguez F, Guerne Y, Vincent JD, Richard P. Release of oxytocin
35
36 1097 within the supraoptic nucleus during the milk ejection reflex in rats. *Exp Brain Res.* 1989; **76**(3): 593-
37
38 1098 602.
39
40 1099 159. Freund-Mercier MJ, Richard P. Electrophysiological evidence for facilitatory control of
41
42 1100 oxytocin neurones by oxytocin during suckling in the rat. *J Physiol.* 1984; **352**447-66.
43
44 1101 160. Neumann I, Douglas AJ, Pittman QJ, Russell JA, Landgraf R. Oxytocin released within the
45
46 1102 supraoptic nucleus of the rat brain by positive feedback action is involved in parturition-related
47
48 1103 events. *J Neuroendocrinol.* 1996; **8**(3): 227-33.
49
50 1104 161. Israel JM, Le Masson G, Theodosis DT, Poulain DA. Glutamatergic input governs periodicity
51
52 1105 and synchronization of bursting activity in oxytocin neurons in hypothalamic organotypic cultures.
53
54 1106 *Eur J Neurosci.* 2003; **17**(12): 2619-29.
55
56
57
58
59
60

- 1107 162. Popescu IR, Buraei Z, Haam J, Weng FJ, Tasker JG. Lactation induces increased IPSC
- 1108 bursting in oxytocinergic neurons. *Physiol Rep*. 2019; **7**(8): e14047.
- 1109 163. Teruyama R, Armstrong WE. Changes in the active membrane properties of rat supraoptic
- 1110 neurones during pregnancy and lactation. *J Neuroendocrinol*. 2002; **14**(12): 933-44.
- 1111 164. Wang YF, Hatton GI. Burst firing of oxytocin neurons in male rat hypothalamic slices. *Brain*
- 1112 *Res*. 2005; **1032**(1-2): 36-43.
- 1113 165. Herbison AE, Voisin DL, Douglas AJ, Chapman C. Profile of monoamine and excitatory
- 1114 amino acid release in rat supraoptic nucleus over parturition. *Endocrinology*. 1997; **138**(1): 33-40.
- 1115 166. Moos F, Richard P. Paraventricular and supraoptic bursting oxytocin cells in rat are locally
- 1116 regulated by oxytocin and functionally related. *J Physiol*. 1989; **408**1-18.
- 1117 167. Hatton GI. Astroglial modulation of neurotransmitter/peptide release from the
- 1118 neurohypophysis: present status. *J Chem Neuroanat*. 1999; **16**(3): 203-21.
- 1119 168. Perlmutter LS, Tweedle CD, Hatton GI. Neuronal/glia plasticity in the supraoptic dendritic
- 1120 zone: dendritic bundling and double synapse formation at parturition. *Neuroscience*. 1984; **13**(3): 769-
- 1121 79.
- 1122 169. Moos F, Marganec A, Fontanaud P, Guillou-Duvoid A, Alonso G. Synchronization of
- 1123 oxytocin neurons in suckled rats: possible role of bilateral innervation of hypothalamic supraoptic
- 1124 nuclei by single medullary neurons. *Eur J Neurosci*. 2004; **20**(1): 66-78.
- 1125 170. Wang YF, Negoro H, Higuchi T. Lesions of hypothalamic mammillary body desynchronise
- 1126 milk-ejection bursts of rat bilateral supraoptic oxytocin neurones. *J Neuroendocrinol*. 2012; **24**: 767-75.
- 1127 171. Boudaba C, Di S, Tasker JG. Presynaptic noradrenergic regulation of glutamate inputs to
- 1128 hypothalamic magnocellular neurones. *J Neuroendocrinol*. 2003; **15**(8): 803-10.
- 1129 172. Roland BL, Sawchenko PE. Local origins of some GABAergic projections to the
- 1130 paraventricular and supraoptic nuclei of the hypothalamus in the rat. *J Comp Neurol*. 1993; **332**(1):
- 1131 123-43.
- 1132 173. Boudaba C, Tasker JG. Internuclear coupling of hypothalamic magnocellular nuclei by
- 1133 glutamate synaptic circuits. *Am J Physiol Regul Integr Comp Physiol*. 2006; **291**(1): R102-R11.

- 1134 174. Sumner BE, Coombes JE, Pumford KM, Russell JA. Opioid receptor subtypes in the
1135 supraoptic nucleus and posterior pituitary gland of morphine-tolerant rats. *Neuroscience*. 1990; **37**(3):
1136 635-45.
- 1137 175. Smith MJ, Wise PM. Localization of kappa opioid receptors in oxytocin magnocellular
1138 neurons in the paraventricular and supraoptic nuclei. *Brain Res*. 2001; **898**(1): 162-5.
- 1139 176. Kim JS, Brown CH, Anderson GM. Anti-opioid effects of RFRP-3 on magnocellular neuron
1140 activity in morphine-naïve and morphine-treated female rats. *Endocrinology*. 2016; **157**(10): 4003-11.
- 1141 177. Martin R, Moll U, Voigt KH. An attempt to characterize by immunocytochemical methods
1142 the enkephalin-like material in oxytocin endings of the rat neurohypophysis. *Life Sci*. 1983; **33 Suppl**
1143 **169**.
- 1144 178. Eriksson M, Ceccatelli S, Uvnas-Moberg K, Iadarola M, Hokfelt T. Expression of Fos-related
1145 antigens, oxytocin, dynorphin and galanin in the paraventricular and supraoptic nuclei of lactating
1146 rats. *Neuroendocrinology*. 1996; **63**(4): 356-67.
- 1147 179. Brown CH, Murphy NP, Munro G, Ludwig M, Bull PM, Leng G, Russell JA. Interruption of
1148 central noradrenergic pathways and morphine withdrawal excitation of oxytocin neurones in the rat. *J*
1149 *Physiol*. 1998; **507**(Pt 3): 831-42.
- 1150 180. Douglas AJ, Neumann I, Meeren HK, Leng G, Johnstone LE, Munro G, Russell JA. Central
1151 endogenous opioid inhibition of supraoptic oxytocin neurons in pregnant rats. *J Neurosci*. 1995; **15**(7
1152 Pt 1): 5049-57.
- 1153 181. Douglas AJ, Bicknell RJ, Leng G, Russell JA, Meddle SL. Beta-endorphin cells in the arcuate
1154 nucleus: Projections to the supraoptic nucleus and changes in expression during pregnancy and
1155 parturition. *J Neuroendocrinol*. 2002; **14**(10): 768-77.
- 1156 182. Zhao BG, Chapman C, Bicknell RJ. Functional kappa-opioid receptors on oxytocin and
1157 vasopressin nerve terminals isolated from the rat neurohypophysis. *Brain Res*. 1988; **462**(1): 62-6.
- 1158 183. Leng G, Dye S, Bicknell RJ. Kappa-opioid restraint of oxytocin secretion: plasticity through
1159 pregnancy. *Neuroendocrinology*. 1997; **66**(6): 378-83.

- 1160 184. Douglas AJ, Bicknell RJ. Oxytocin nerve terminal desensitization to kappa-opioids at the end
1161 of pregnancy. *Ann N Y Acad Sci.* 1993; **689**:589-92.
- 1162 185. Douglas AJ, Dye S, Leng G, Russell JA, Bicknell RJ. Endogenous opioid regulation of
1163 oxytocin secretion through pregnancy in the rat. *J Neuroendocrinol.* 1993; **5**(3): 307-14.
- 1164 186. Srisawat R, Ludwig M, Bull PM, Douglas AJ, Russell JA, Leng G. Nitric oxide and the
1165 oxytocin system in pregnancy. *J Neurosci.* 2000; **20**(17): 6721-37.
- 1166 187. Bull PM, Ludwig M, Blackburn-Munro GJ, Delgado-Cohen H, Brown CH, Russell JA. The
1167 role of nitric oxide in morphine dependence and withdrawal excitation of rat oxytocin neurons. *Eur J*
1168 *Neurosci.* 2003; **18**(9): 2545-51.
- 1169 188. Doi N, Brown CH, Cohen HD, Leng G, Russell JA. Effects of the endogenous opioid peptide,
1170 endomorphin 1, on supraoptic nucleus oxytocin and vasopressin neurones *in vivo* and *in vitro* *Br J*
1171 *Pharmacol.* 2001; **132**(5): 1136-44.
- 1172 189. Brown CH, Russell JA. Cellular mechanisms underlying neuronal excitability during
1173 morphine withdrawal in physical dependence: lessons from the magnocellular oxytocin system.
1174 *Stress.* 2004; **7**(2): 97-107.
- 1175 190. Brown CH, Munro G, Johnstone LE, Robson AC, Landgraf R, Russell JA. Oxytocin neurone
1176 autoexcitation during morphine withdrawal in anaesthetized rats. *Neuroreport.* 1997; **8**(4): 951-5.
- 1177 191. Pinol RA, Jameson H, Popratiloff A, Lee NH, Mendelowitz D. Visualization of oxytocin
1178 release that mediates paired pulse facilitation in hypothalamic pathways to brainstem autonomic
1179 neurons. *PLoS One.* 2014; **9**(11): e112138.
- 1180 192. Roy RK, Augustine RA, Brown CH, Schwenke DO. Activation of oxytocin neurons in the
1181 paraventricular nucleus drives cardiac sympathetic nerve activation following myocardial infarction in
1182 rats. *Commun Biol.* 2018; **1**:160.
- 1183 193. Son SJ, Filosa JA, Potapenko ES, Biancardi VC, Zheng H, Patel KP, Tobin VA, Ludwig M,
1184 Stern JE. Dendritic peptide release mediates interpopulation crosstalk between neurosecretory and
1185 preautonomic networks. *Neuron.* 2013; **78**(6): 1036-49.

- 1
- 2
- 3 1186 194. Zaelzer C, Gizowski C, Salmon CK, Murai KK, Bourque CW. Detection of activity-
- 4
- 5 1187 dependent vasopressin release from neuronal dendrites and axon terminals using sniffer cells. *J*
- 6
- 7 1188 *Neurophysiol.* 2018; **120**(3): 1386-96.
- 8
- 9 1189 195. Theodosios DT, Poulain DA. Evidence that oxytocin-secreting neurones are involved in the
- 10
- 11 1190 ultrastructural reorganisation of the rat supraoptic nucleus apparent at lactation. *Cell Tissue Res.* 1984;
- 12
- 13 1191 **235**(1): 217-9.
- 14
- 15 1192 196. Tweedle CD, Hatton GI. Ultrastructural changes in rat hypothalamic neurosecretory cells and
- 16
- 17 1193 their associated glia during minimal dehydration and rehydration. *Cell Tissue Res.* 1977; **181**(1): 59-
- 18
- 19 1194 72.
- 20
- 21 1195 197. Piet R, Vargova L, Sykova E, Poulain DA, Oliet SH. Physiological contribution of the
- 22
- 23 1196 astrocytic environment of neurons to intersynaptic crosstalk. *Proc Natl Acad Sci U S A.* 2004; **101**(7):
- 24
- 25 1197 2151-5.
- 26
- 27 1198 198. Strack AM, Sawyer WB, Hughes JH, Platt KB, Loewy AD. A general pattern of CNS
- 28
- 29 1199 innervation of the sympathetic outflow demonstrated by transneuronal pseudorabies viral infections.
- 30
- 31 1200 *Brain Res.* 1989; **491**(1): 156-62.
- 32
- 33 1201 199. Bourque CW. Central mechanisms of osmosensation and systemic osmoregulation. *Nat Rev*
- 34
- 35 1202 *Neurosci.* 2008; **9**(7): 519-31.
- 36
- 37 1203 200. Toney GM, Chen QH, Cato MJ, Stocker SD. Central osmotic regulation of sympathetic nerve
- 38
- 39 1204 activity. *Acta Physiol Scand.* 2003; **177**(1): 43-55.
- 40
- 41 1205 201. Gabor A, Leenen FH. Central neuromodulatory pathways regulating sympathetic activity in
- 42
- 43 1206 hypertension. *J Appl Physiol* 2012; **113**(8): 1294-303.
- 44
- 45 1207 202. Han SY, Gray E, Hughes G, Brown CH, Schwenke DO. Increased sympathetic drive during
- 46
- 47 1208 the onset of hypertension in conscious Cyp1a1-Ren2 rats. *Pflugers Arch Eur J Physiol.* 2014; **466**459-
- 48
- 49 1209 66.
- 50
- 51 1210 203. Jardine DL, Charles CJ, Ashton RK, Bennett SI, Whitehead M, Frampton CM, Nicholls MG.
- 52
- 53 1211 Increased cardiac sympathetic nerve activity following acute myocardial infarction in a sheep model.
- 54
- 55 1212 *J Physiol.* 2005; **565**(Pt 1): 325-33.
- 56
- 57
- 58
- 59
- 60

- 1213 204. Leenen FH. Brain mechanisms contributing to sympathetic hyperactivity and heart failure.
1214 *Circ Res.* 2007; **101**(3): 221-3.
- 1215 205. Biancardi VC, Son SJ, Sonner PM, Zheng H, Patel KP, Stern JE. Contribution of central
1216 nervous system endothelial nitric oxide synthase to neurohumoral activation in heart failure rats.
1217 *Hypertension.* 2011; **58**(3): 454-63.
- 1218 206. Cohn JN, Levine TB, Olivari MT, Garberg V, Lura D, Francis GS, Simon AB, Rector T.
1219 Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. *N*
1220 *Engl J Med.* 1984; **311**(13): 819-23.
- 1221 207. Han SY, Bouwer GT, Seymour AJ, Korpak AK, Schwenke DO, Brown CH. Induction of
1222 hypertension blunts baroreflex inhibition of vasopressin neurons in the rat. *Eur J Neurosci.* 2015;
1223 **42**(9): 2690-8.
- 1224 208. Korpak AK, Han SY, Schwenke DO, Brown CH. A switch from GABA inhibition to
1225 excitation of vasopressin neurons exacerbates the development angiotensin II-dependent
1226 hypertension. *J Neuroendocrinol.* 2018; **30** e12564.
- 1227 209. Roy RK, Augustine RA, Brown CH, Schwenke DO. Acute myocardial infarction activates
1228 magnocellular vasopressin and oxytocin neurons. *J Neuroendocrinol.* 2019; **31**e12808.
- 1229 210. Potapenko ES, Biancardi VC, Zhou Y, Stern JE. Altered astrocyte glutamate transporter
1230 regulation of hypothalamic neurosecretory neurons in heart failure rats. *Am J Physiol Regul Integr*
1231 *Comp Physiol.* 2012; **303**(3): R291-300.
- 1232 211. Stern JE, Potapenko ES. Enhanced NMDA receptor-mediated intracellular calcium signaling
1233 in magnocellular neurosecretory neurons in heart failure rats. *Am J Physiol Regul Integr Comp Physiol.*
1234 2013; **305**(4): R414-22.
- 1235 212. Goldsmith SR, Francis GS, Cowley AW, Jr., Levine TB, Cohn JN. Increased plasma arginine
1236 vasopressin levels in patients with congestive heart failure. *J Am Coll Cardiol.* 1983; **1**(6): 1385-90.
- 1237 213. Goldsmith SR, Gheorghiade M. Vasopressin antagonism in heart failure. *J Am Coll Cardiol.*
1238 2005; **46**(10): 1785-91.

- 1239 214. Murakami K, Akana S, Dallman MF, Ganong WF. Correlation between the stress-induced
- 1240 transient increase in corticotropin-releasing hormone content of the median eminence of the
- 1241 hypothalamus and adrenocorticotrophic hormone secretion. *Neuroendocrinology*. 1989; **49**(3): 233-41.
- 1242 215. Dabrowska J, Hazra R, Ahern TH, Guo JD, McDonald AJ, Mascagni F, Muller JF, Young LJ,
- 1243 Rainnie DG. Neuroanatomical evidence for reciprocal regulation of the corticotrophin-releasing factor
- 1244 and oxytocin systems in the hypothalamus and the bed nucleus of the stria terminalis of the rat:
- 1245 Implications for balancing stress and affect. *Psychoneuroendocrinology*. 2011; **36**(9): 1312-26.
- 1246 216. Jamieson BB, Nair BB, Iremonger KJ. Regulation of hypothalamic corticotropin-releasing
- 1247 hormone neurone excitability by oxytocin. *J Neuroendocrinol*. 2017; **29**(11): e12532.
- 1248 217. Attwell D, Buchan AM, Charkpak S, Lauritzen M, Macvicar BA, Newman EA. Glial and
- 1249 neuronal control of brain blood flow. *Nature*. 2010; **468**(7321): 232-43.
- 1250 218. Du W, Stern JE, Filosa JA. Neuronal-derived nitric oxide and somatodendritically released
- 1251 vasopressin regulate neurovascular coupling in the rat hypothalamic supraoptic nucleus. *J Neurosci*.
- 1252 2015; **35**(13): 5330-41.
- 1253 219. Leng G, Sabatier N. Oxytocin - The sweet hormone? *Trends Endocrinol Metab*. 2017; **28**(5):
- 1254 365-76.
- 1255 220. Sabatier N, Caquineau C, Douglas AJ, Leng G. Oxytocin released from magnocellular
- 1256 dendrites: a potential modulator of alpha-melanocyte-stimulating hormone behavioral actions? *Ann N*
- 1257 *Y Acad Sci*. 2003; **994**218-24.
- 1258 221. Klockars OA, Waas JR, Klockars A, Levine AS, Olszewski PK. Neural basis of ventromedial
- 1259 hypothalamic oxytocin-driven decrease in appetite. *Neuroscience*. 2017; **366**54-61.
- 1260 222. Patisaul HB, Scordalakes EM, Young LJ, Rissman EF. Oxytocin, but not oxytocin receptor, is
- 1261 regulated by oestrogen receptor beta in the female mouse hypothalamus. *J Neuroendocrinol*. 2003;
- 1262 **15**(8): 787-93.
- 1263 223. Manning M, Misicka A, Olma A, Bankowski K, Stoev S, Chini B, Durroux T, Mouillac B,
- 1264 Corbani M, Guillon G. Oxytocin and vasopressin agonists and antagonists as research tools and
- 1265 potential therapeutics. *J Neuroendocrinol*. 2012; **24**(4): 609-28.

- 1266 224. Chen C, Jiang Z, Fu X, Yu D, Huang H, Tasker JG. Astrocytes amplify neuronal dendritic
1267 volume transmission stimulated by norepinephrine. *Cell Rep.* 2019; **29**(13): 4349-61.e4.
- 1268 225. Takayanagi Y, Yoshida M, Takashima A, Takanami K, Yoshida S, Nishimori K, Nishijima I,
1269 Sakamoto H, Yamagata T, Onaka T. Activation of supraoptic oxytocin neurons by secretin facilitates
1270 social recognition. *Biol Psychiatry.* 2017; **81**(3): 243-51.
- 1271 226. Ludwig M, Leng G. Dendritic peptide release and peptide-dependent behaviours. *Nat Rev*
1272 *Neurosci.* 2006; **7**(2): 126-36.
- 1273 227. Chini B, Verhage M, Grinevich V. The Action Radius of Oxytocin Release in the Mammalian
1274 CNS: From Single Vesicles to Behavior. *Trends Pharmacol Sci.* 2017; **38**(11): 982-91.
- 1275 228. Mens WB, Witter A, van Wimersma Greidanus TB. Penetration of neurohypophyseal
1276 hormones from plasma into cerebrospinal fluid (CSF): half-times of disappearance of these
1277 neuropeptides from CSF. *Brain Res.* 1983; **262**(1): 143-9.
- 1278 229. Stark H, Burbach JP, Van der Kleij AA, De Wied D. In vivo conversion of vasopressin after
1279 microinjection into limbic brain areas of rats. *Peptides.* 1989; **10**(4): 717-20.
- 1280 230. Xia X, Lessmann V, Martin TF. Imaging of evoked dense-core-vesicle exocytosis in
1281 hippocampal neurons reveals long latencies and kiss-and-run fusion events. *J Cell Sci.* 2009; **122**(Pt
1282 1): 75-82.
- 1283 231. Engelmann M, Ebner K, Landgraf R, Wotjak CT. Effects of Morris water maze testing on the
1284 neuroendocrine stress response and intrahypothalamic release of vasopressin and oxytocin in the rat.
1285 *Horm Behav.* 2006; **50**(3): 496-501.
- 1286 232. Ebner K, Wotjak CT, Landgraf R, Engelmann M. Forced swimming triggers vasopressin
1287 release within the amygdala to modulate stress-coping strategies in rats. *Eur J Neurosci.* 2002; **15**(2):
1288 384.
- 1289 233. Ebner K, Wotjak CT, Landgraf R, Engelmann M. A single social defeat experience
1290 selectively stimulates the release of oxytocin, but not vasopressin, within the septal brain area of male
1291 rats. *Brain Res.* 2000; **872**(1-2): 87.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1292 234. Engelmann M, Ebner K, Landgraf R, Wotjak CT. Swim stress triggers the release of
1293 vasopressin within the suprachiasmatic nucleus of male rats. *Brain Res.* 1998; **792**(2): 343.
1294 235. Ebner K, Wotjak CT, Holsboer F, Landgraf R, Engelmann M. Vasopressin released within
1295 the septal brain area during swim stress modulates the behavioural stress response in rats.
1296 *Eur J Neurosci.* 1999; **11**(3): 997.
1297 236. Landgraf R, Ludwig M. Vasopressin release within the supraoptic and paraventricular nuclei
1298 of the rat brain: osmotic stimulation via microdialysis. *Brain Res.* 1991; **558**(2): 191.
1299 237. Althammer F, Grinevich V. Diversity of oxytocin neurons: beyond magno- and parvocellular
1300 cell types? *J Neuroendocrinol.* 2017; **12**10.1111/jne.12549.
1301 238. Ludwig M, Sabatier N, Dayanithi G, Russell JA, Leng G. The active role of dendrites in the
1302 regulation of magnocellular neurosecretory cell behavior. *Prog Brain Res.* 2002; **139**247.
1303 239. Stern JE. Neuroendocrine-autonomic integration in the paraventricular nucleus: novel roles
1304 for dendritically released neuropeptides. *J Neuroendocrinol.* 2015; **27**(6): 487-97.

Figure legends

Figure 1. Magnocellular neurosecretory cells. Magnocellular neurosecretory cells (MNCs) of the hypothalamic supraoptic nucleus and paraventricular nucleus each possess 1 – 3 dendrites and project a single axon to the posterior pituitary gland where they secrete either oxytocin or vasopressin into the circulation. Some MNC axons project axon collaterals to other brains areas.

Figure 2. Mechanisms of somato-dendritic release of oxytocin from magnocellular neurosecretory cells. Neuropeptides are synthesized and packaged in the soma and stored in dendrites in a reserve pool containing large numbers of large dense core vesicles (LDCVs). Depolarization-induced calcium entry through voltage-gated calcium channels (VGCCs) stimulates peptide release by exocytosis of LDCVs. This requires the depolymerization of F-actin to G-actin. Furthermore, the stimulation of G-protein coupled receptors (GPR), such as the oxytocin receptor, stimulates the mobilization of Ca^{2+} from IP₃-dependent intracellular stores of the rough endoplasmic reticulum (ER) and an increase in the number of LDCVs at the plasma membrane, thus priming the exocytosis machinery for subsequent activity-dependent release. Although some members of the SNARE family are detectable by immunocytochemistry, there appears to be a lack of VAMP, SNAP-25 and synaptotagmin-1 in the somata and dendrites, with their function presumably being replaced by other SNARE proteins.

Figure 3. Autocrine modulation of burst firing in oxytocin magnocellular neurosecretory cells. Cervical stretch during birth activates stretch receptors to activate A2 noradrenergic neurones in the nucleus tractus solitarius (NTS) that, in turn, activates somato-dendritic oxytocin secretion from oxytocin magnocellular neurosecretory cells (MNCs). Oxytocin feeds back on oxytocin MNCs to increase excitability. Oxytocin also increases noradrenaline secretion within the SON to establishes a local positive feedback loop that reinforces oxytocin MNC excitation and promotes oxytocin secretion into the circulation to trigger uterine contractions during birth.

Figure 4. Ghrelin stimulation of somato-dendritic vasopressin secretion. Ghrelin activation of growth hormone secretagogue receptors (GHSR) on vasopressin magnocellular neurosecretory cells (MNCs) induces somato-dendritic vasopressin secretion, which activates V_{1a} receptors ($V_{1a}Rs$) on neighbouring astrocytes to increase intracellular calcium. Increased astrocytic calcium triggers release of the gliotransmitter, ATP, which activates ionotropic P2X receptors on GABA interneurons that project back to vasopressin MNCs.

Figure 5. Autocrine modulation of vasopressin magnocellular neurosecretory cell activity. Vasopressin magnocellular neurosecretory cells (MNCs) secrete vasopressin, ATP and dynorphin (and other transmitters) from their somata and dendrites. Endogenous AVP (2) inhibits spike discharge throughout bursts via inhibition of EPSC amplitude. Endogenous ATP is rapidly converted to adenosine (3), which enhances the medium afterhyperpolarisation (mAHP) amplitude over the first few seconds of bursts to contribute to spike frequency adaptation. Endogenous dynorphin (4) inhibition of the afterdepolarisation (ADP) increases progressively over the course of bursts, eventually resulting in burst termination. Combined, these autocrine feedback effects of somato-dendritic vasopressin and co-secreted transmitters shape phasic activity for efficient secretion of vasopressin into the circulation from the posterior pituitary gland.

Figure 6. Endocannabinoid modulation of excitatory and inhibitory synapses on MNCs. Oxytocin activation of autocrine oxytocin receptors (OTR) on oxytocin neurons leads to a tonic basal release of the endocannabinoid anandamide (AEA) at GABA synapses, which tonically suppresses synaptic inhibitory input to oxytocin neurons by activating presynaptic CB1 receptors. Depolarization (e.g., via action potential generation) or corticosteroid (Cort) exposure (e.g., during stress) leads to a calcium-dependent release of the other main endocannabinoid, 2-arachidonoylglycerol (2-AG), at glutamate synapses, which suppresses synaptic excitation of both oxytocin and vasopressin MNCs by activating presynaptic CB1 receptors. Glial retraction induced by salt loading allows the 2-AG released at glutamate

synapses to spill over onto GABA synapses and suppress synaptic inhibition via CB1 receptor activation. Tonic AEA occupation of CB1 receptors at GABA synapses is non-saturating, allowing additional suppression of GABA release following phasic 2-AG release and synaptic spillover.

Figure 7. Paracrine actions of somato-dendritic vasopressin secretion. Activation of neurosecretory vasopressin magnocellular neurosecretory cells (MNCs) (1) triggers action potential firing (2) to release vasopressin into the circulation from the posterior pituitary gland (3). In parallel, action potentials back-propagate into the dendrites (4) to trigger somato-dendritic vasopressin secretion (5). In addition to autocrine feedback inhibition of vasopressin MNC activity via V_{1a} receptors ($V_{1a}Rs$) (6), somato-dendritic vasopressin diffuses through the extracellular space to bind to $V_{1a}Rs$ on presympathetic paraventricular nucleus neurones (7) to increase action potential firing (8) and therefore increase sympathetic outflow to peripheral organs. Somato-dendritic vasopressin also activates $V_{1a}Rs$ on local blood vessels (9) to cause vasoconstriction, which is predicted to inhibit vasopressin MNCs at a population level (10) by restricting the availability of oxygen and nutrients. Hence, somato-dendritic vasopressin secretion coordinates neurohumoral responses to (patho)physiological activation (11).




Figure 1. Magnocellular neurosecretory cells. Magnocellular neurosecretory cells (MNCs) of the hypothalamic supraoptic nucleus and paraventricular nucleus each possess 1 – 3 dendrites and project a single axon to the posterior pituitary gland where they secrete either oxytocin or vasopressin into the circulation. Some MNC axons project axon collaterals to other brains areas.

185x134mm (300 x 300 DPI)





Figure 2. Mechanisms of somato-dendritic release of oxytocin from magnocellular neurosecretory cells.

Neuropeptides are synthesized and packaged in the soma and stored in dendrites in a reserve pool containing large numbers of large dense core vesicles (LDCVs). Depolarization-induced calcium entry through voltage-gated calcium channels (VGCCs) stimulates peptide release by exocytosis of LDCVs. This requires the depolymerization of F-actin to G-actin. Furthermore, the stimulation of G-protein coupled receptors (GPR), such as the oxytocin receptor, stimulates the mobilization of Ca^{2+} from IP_3 -dependent intracellular stores of the rough endoplasmic reticulum (ER) and an increase in the number of LDCVs at the plasma membrane, thus priming the exocytosis machinery for subsequent activity-dependent release. Although some members of the SNARE family are detectable by immunocytochemistry, there appears to be a lack of VAMP, SNAP-25 and synaptotagmin-1 in the somata and dendrites, with their function presumably being replaced by other SNARE proteins.

181x163mm (300 x 300 DPI)



179x181mm (300 x 300 DPI)





Figure 4. Ghrelin stimulation of somato-dendritic vasopressin secretion. Ghrelin activation of growth hormone secretagogue receptors (GHSR) on vasopressin magnocellular neurosecretory cells (MNCs) induces somato-dendritic vasopressin secretion, which activates V1a receptors (V1aRs) on neighbouring astrocytes to increase intracellular calcium. Increased astrocytic calcium triggers release of the gliotransmitter, ATP, which activates ionotropic P2X receptors on GABA interneurons that project back to vasopressin MNCs.

179x218mm (300 x 300 DPI)



179x246mm (300 x 300 DPI)




Figure 6. Endocannabinoid modulation of excitatory and inhibitory synapses on MNCs. Oxytocin activation of autocrine oxytocin receptors (OTR) on oxytocin neurons leads to a tonic basal release of the endocannabinoid anandamide (AEA) at GABA synapses, which tonically suppresses synaptic inhibitory input to oxytocin neurons by activating presynaptic CB1 receptors. Depolarization (e.g., via action potential generation) or corticosteroid (Cort) exposure (e.g., during stress) leads to a calcium-dependent release of the other main endocannabinoid, 2-arachidonoylglycerol (2-AG), at glutamate synapses, which suppresses synaptic excitation of both oxytocin and vasopressin MNCs by activating presynaptic CB1 receptors. Glial retraction induced by salt loading allows the 2-AG released at glutamate synapses to spill over onto GABA synapses and suppress synaptic inhibition via CB1 receptor activation. Tonic AEA occupation of CB1 receptors at GABA synapses is non-saturating, allowing additional suppression of GABA release following phasic 2-AG release and synaptic spillover.

179x154mm (300 x 300 DPI)




Figure 7. Paracrine actions of somato-dendritic vasopressin secretion. Activation of neurosecretory vasopressin magnocellular neurosecretory cells (MNCs) (1) triggers action potential firing (2) to release vasopressin into the circulation from the posterior pituitary gland (3). In parallel, action potentials back-propagate into the dendrites (4) to trigger somato-dendritic vasopressin secretion (5). In addition to autocrine feedback inhibition of vasopressin MNC activity via V1a receptors (V1aRs) (6), somato-dendritic vasopressin diffuses through the extracellular space to bind to V1aRs on presympathetic paraventricular nucleus neurones (7) to increase action potential firing (8) and therefore increase sympathetic outflow to peripheral organs. Somato-dendritic vasopressin also activates V1aRs on local blood vessels (9) to cause vasoconstriction, which is predicted to inhibit vasopressin MNCs at a population level (10) by restricting the availability of oxygen and nutrients. Hence, somato-dendritic vasopressin secretion coordinates neurohumoral responses to (patho)physiological activation (11).

179x183mm (300 x 300 DPI)